

This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

+ *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.

+ *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.

+ *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.

+ *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at `http://books.google.com/`

S .33.1

HARVARD COLLEGE
LIBRARY



FROM THE LIBRARY OF

WILLIAM LAMBERT RICHARDSON

A.B. 1864, M.D. 1867

OF BOSTON

THE MUSHROOM BOOK



Amanitopsis parcivolvata, Pk.

See page 55.

PAGE

Genus Psathyra 82
Genus Psilocybe 82
Genus Pilosace 82
Genus Deconica 82
Genus Chitonia 83

3. RUSTY-SPORED SERIES

Genus Pholiota 83
Fat Pholiota 83
Early Pholiota 84
Pholiota aggericola 84
Genus Cortinarius 85
Cortinarius alboviolaceus 86

4. PINK-SPORED SERIES

Genus Pluteus 87
Fawn-coloured Pluteus 87
Genus Entoloma 88
Genus Eccilia 88
Genus Volvaria 88
Genus Clitopilus 88

5. BLACK-SPORED SERIES

Genus Coprinus 89
Ink Caps 89
Shaggy-mane; Horsetail; Maned Agaric . 90
Inky Coprinus 91
Glistening Coprinus 91
Genus Gomphidius 92
Genus Psathyrella 92
Genus Panæolus 92

6. FUNGI WITH MILKY JUICE

Genus Lactarius 92
Peppery Lactarius 92
Lactarius ligniotus 93

CHAPTER VI: FUNGI WITH TEETH—HYDNACEÆ . . 94

Genus Hydnum 95
Spreading Hydnum 95
White Hydnum 95

Copyright, 1901, 1904, by
Doubleday, Page & Company

Norwood Press:
Berwick & Smith Co., Norwood, Mass., U.S.A.

THE MUSHROOM BOOK. A POPULAR GUIDE TO THE IDENTIFICATION AND STUDY OF OUR COMMONER FUNGI, WITH SPECIAL EMPHASIS ON THE EDIBLE VARIETIES

BY

NINA L. MARSHALL

WITH MANY ILLUSTRATIONS IN COLOR AND BLACK AND WHITE PHOTOGRAPHED FROM NATURE BY J. A. & H. C. ANDERSON

NEW YORK
DOUBLEDAY, PAGE & COMPANY
1905

NH 568.4(10)
S 8000.1(10)

HARVARD COLLEGE LIBRARY
BEQUEST OF
DR. WILLIAM L. RICHARDSON
FEBRUARY 24, 1933

Copyright, 1901, 1904, by
Doubleday, Page & Company

HARVARD
UNIVERSITY
LIBRARY
JUL 22 1966

Norwood Press:
Berwick & Smith Co., Norwood, Mass., U.S.A.



THE MUSHROOM BOOK

COPRINUS COMATUS
Courtesy of Agricultural Experiment Station, Cornell.
See page 90

PAGE

The preparation of rough-dried plants for the herbarium 149
Mounting 149
Sections 150
Poisoning herbarium specimens 150
CHAPTER XV : FUNGI FOR THE TABLE 151
Cautions for the inexperienced 151
The food value of fungi 152
To keep mushrooms temporarily 153
To prepare the edible agarics for cooking 153
To toast agarics 153
To bake agarics 153
To broil agarics 153
Mushrooms stewed 153
To prepare russulas 154
To prepare fungi with milky juice 154
Lactarius deliciosus 154
Lactarius volemus 154
To prepare amanitas 154
To prepare chanterelles 154
To fry chanterelles 154
To prepare coprini—ink caps 154
To prepare boleti 154
To prepare Hydnum repandum 154
To prepare morels 154
To prepare beefsteak fungus 155
For salad 155
Minced 155
To prepare gyromitras 155
To prepare woody pore-bearing fungi—Polyporæ . 155
To prepare clavarias and branched hydnums . . 155
To cook clavarias 155
To prepare puffballs 155
To cook puffballs 155
To cook the giant puffball 155
Puffball salad 155
LIST OF AUTHORITIES CONSULTED 157
ABBREVIATIONS OF NAMES OF BOTANISTS WITH EXPLANATIONS . 159
INDEX AND GLOSSARY 161

LIST OF PLATES

PLATE		FACING PAGE
I.	Amanitopsis parcivolvata, Pk. (see p. 55)	*Frontispiece*
II.	Death Cup; Poison Amanita (Amanita phalloides, Fr.)	48
III.	Fly Amanita (Amanita muscaria, L.), poisonous	49
IV.	Orange Amanita (Amanita Cæsarea, Scop.), edible	50
V.	Strangled Amanitopsis (Amanitopsis strangulata, Fr. Roze), edible	53
VI.	Sheathed Amanitopsis (Amanitopsis vaginata, Roze), edible	54
VII.	Helmet Mycena (Mycena galericulata, Scop.), edible	55
VIII.	Scaly Lentinus (Lentinus lepideus, Fr.), edible	56
IX.	Oyster Mushroom (Pleurotus ostreatus, Fr.), edible (see p. 57)	58
X.	Golden-flesh Boletus (Boletus chrysenteron, Fr.) (see p. 106)	60
	Vermilion Hygrophorus (Hygrophorus miniatus, Fr.), edible	60
	Chantarelle Hygrophorus (Hygrophorus cantharellus, Schw.), edible (see p. 59)	60
XI.	Fat Pholiota (Pholiota adiposa, Fr.), edible (see p. 83)	61
	Honey-coloured Armillaria (Armillaria mellea, Vahl.), edible	61
XII.	Grainy Lepiota (Lepiota granosa, Morg.),	63
XIII.	Smooth Lepiota (Lepiota naucinoides, Pk.), edible	64
	Parasol Mushroom (Lepiota procera, Scop.), edible (see p. 63)	64
XIV.	Lepiota Friesii, Lasch	65
	Cortinarius alboviolaceus, Fr. (see p. 86)	65

PLATE		FACING PAGE
XV.	Spotted Collybia (Collybia maculata, A. and S.)	66
	Broad-gilled Collybia (Collybia platyphylla, Fr.)	66
XVI.	Collybia familia, Pk. (edible)	67
	Clitocybe laccata, Scop., edible (see p. 70) .	67
XVII.	Emetic Russula (Russula emetica, Schaeff.), (dangerous)	68
XVIII.	Green Russula (Russula virescens, Fr.), edible .	69
XIX.	Deceiving Clitocybe (Clitocybe illudens, Schw.)	70
XX.	Clitocybe virens, Scop. (edible)	71
XXI.	Masked Tricholoma (Tricholoma personatum, Fr.; var. bulbosum, Pk.), edible . . .	72
XXII.	Pholiota aggericola, Peck (see p. 84) . . .	73
	Craterellus cantharellus, Schw. (edible) . .	73
XXIII.	Agaricus campestris, L., edible (see p. 76) .	74
	Agaricus abruptus, Pk. (edible)	74
XXIV.	Field Mushroom (Agaricus campestris, L.), edible (uncultivated)	75
XXV.	Rodman's Mushroom (Agaricus Rodmani, Pk.), young	76
	Rodman's Mushroom (mature)	76
XXVI.	Agaricus abruptus, Pk. (edible)	77
XXVII.	Perplexing Hypholoma (Hypholoma perplexum, Pk.), harmless	78
XXVIII.	Uncertain Hypholoma (Hypholoma incertum, Pk.), edible . . . . . ,	80
XXIX.	Brick Top (Hypholoma sublateritium, Schaeff.), edible	81
XXX.	Ivory Hygrophorus (Hygrophorus eburneus, Fr.), edible (see p. 61)	84
	Early Pholiota (Pholiota præcox, Pers.), edible .	84
XXXI.	Dog Cortinarius (Cortinarius caninus, Fr.) . .	85
XXXII.	Zoned Cortinarius (Cortinarius armillatus, A. and S., Fr.)	86

PLATE		FACING PAGE
XXXIII.	Fawn-coloured Pluteus (Pluteus cervinus, Schaeff.), edible	87
XXXIV.	Inky Coprinus (Coprinus atramentarius, Fr.; var. silvestris, Pk.), edible (see p. 91) . .	89
XXXV.	Inky Coprinus (Coprinus atramentarius, Fr.), edible	91
XXXVI.	Peppery Lactarius (Lactarius piperatus, Scop.), edible	92
XXXVII.	Mycena hæmatopoda, Pers. (see p. 55) . .	93
	Lactarius ligniotus, Fr.	93
XXXVIII.	Bear's-head (Hydnum caput-ursi, Fr.), edible .	96
XXXIX.	Coral Hydnum (Hydnum coralloides, Scop.), edible	97
XL.	Little Tongue Clavaria (Clavaria ligula, Fr.) .	98
XLI.	Golden Clavaria (Clavaria aurea, Schaeff.), edible	100
XLII.	Clavaria formosa, Pers., edible	101
XLIII.	Cone-like Boletus (Strobilomyces strobilaceus, Berk.)	102
XLIV.	Painted Boletinus (Boletinus pictus, Pk.), edible	103
	Spreading Hydnum (Hydnum repandum, L.), edible (see p. 95)	103
XLV.	Bitter Boletus (Boletus felleus, Bull.; var. obesus, Pk.)	107
XLVI.	Scabrous-stemmed Boletus (Boletus scaber, Fr.; var. niveus, Gill.)	108
XLVII.	Tinder-wood Polyporus (Elfingia fomentaria, L., Fomes fomentarius, Gill., Polyporus fomentarius, Fr.)	110
XLVIII.	Polyporus versicolor, Fr.	112
	Polyporus circinatus, Fr. (see p. 113) . .	112
XLIX.	Lenzites betulina, Fr.	113
L.	Dædalea quercina, L., Pers.	114
LI.	Jew's Ear (Hirneola auricula-Judæ), L., Berk. .	116

PLATE		FACING PAGE
LI.	Xylaria (see p. 136)	116
LII.	Phallus impudicus, L.	119
LIII.	Pear-shaped Puffball (Lycoperdon pyriforme, Schaeff.), edible	125
LIV.	Brain Puffball (Calvatia craniformis, Schw.), edible	126
LV.	Bovistella Ohiensis (Ellis and Morgan), edible (see p. 129)	128
	Cup-shaped Calvatia (Calvatia cyathiformis, Bosc), edible	128
LVI.	Least Earth-star (Geaster minimus, Schw.)	130
	Water-measuring Earth-star (Geaster hygrometricus, Pers.)	130
	Bird's Nest (Cyathus vernicosus, D. C.) (see p. 133)	130
LVII.	Calostoma Ravenelli, Berk., Mass.	132
	Calostoma lutescens, Schw., Burnap	132
	Calostoma cinnabarinum, Desv.	132
	Spathularia velutipes, C. and F. (see p. 138)	132
LVIII.	Flesh-coloured Puffball (Lycoperdon subincarnatum, Pk.), edible (see p. 125)	134
	Young Pear-shaped Puffball (Lycoperdon pyriforme, Schaeff.), edible (see p. 125)	134
	Hard-skinned Puffball (Scleroderma vulgare, Fr.)	134
LIX.	Cordyceps capitala (Holmsk., Lk.), parasitic on Elaphomyces	136
	Lycogola epidendron (see p. 144)	136
	Floccose Chanterelle (Cantharellus floccosus, Schw.), edible (see p. 53)	136
	Mutinus caninus, Huds. (see p. 120)	136
LX.	Slippery Leotia (Leotia lubrica), edible	137
	Jelly-like Tremellodon (Tremellodon gelatinosum)	137
LXI.	Golden Peziza (Peziza aurantia, Pers.), edible	138

PLATE		FACING PAGE
LXI.	Peziza odorata, Pk., edible (see p. 137) . .	138
LXII.	Helvella elastica, Bull. (see p. 142) . . .	140
	Helvella lacunosa, Holm. (see p. 143) . . .	140
	Mitrula vitellina, Sacc., var. irregularis, Pk. . .	140
LXIII.	Delicious Morel (Morchella deliciosa, Fr.), edible .	142
	Polyporus arcularius, Batsch, Fr. (see p. 112) .	142
LXIV.	Bristly Panus (Panus strigosus, B. and C.) (see p. 67)	145

LIST OF ILLUSTRATIONS IN TEXT

	Page
Coprinus comatus, Fr. (old)	i
Coprinus comatus, Fr. (mature)	v
Coprinus comatus, Fr. (young)	viii
Puffball	4, 25
Corollas and honey, attractive to insects	5
Ingenious stamens	5
Pistil of violet	6
Seed-box of iris	6
Pistil of St. Johnswort	6
Seed-box of sacred bean	6
Winged seed of the silver fir	6
Fern with spores (Polypodium vulgare)	6
Grass spikelet	7
Liverwort (Marchantia polymorpha)	7
Bean seed to show embryo	7
Indian pipe	7
Moss (Dicranum scoparium) (natural size)	8
White mould on dead fly	8
Spores borne in delicate membranous sacs	9, 17
Bread mould	9
Spores borne on little spicules	9, 17
Spores as simple cells	11
Spores divided into several cells	11
Mycelial threads	11
Mushroom buttons	11
Young mushroom	11
Puffball	12

List of Illustrations in Text

PAGE

Mushroom to show veil 12
Mushroom to show cap, ring, gills, and stem . . . 12
Mushroom to show veil on stem and cap 12
Section across gill (magnified) 12
A small portion of section of gill (highly magnified) . . 13
Corn smut 13, 22
Leaf rust on Hepatica triloba 13, 22
Pouch-fungus, section to show spores in hollow rind . 14
Section to show gills 14
Section of a Boletus, to show pores 14, 102
Clavaria with spores on spines 14
Section of Hydnum, to show teeth 14
Section of stroma with perithecia (magnified) . . . 18
Perithecium (highly magnified) 18
Outer surface of truffle 18
Section of truffle 18
Fleshy cup-like fungi 19, 20
Club-shaped fungus 19, 21
Conic, convex, and pitted fungus 19–21
Fungus with gyrose furrows 19, 20
Saddle-shaped fungus 19, 20
Fungus with spore surface flat, running down the sides . 21
Fungus with gills 22, 30
Fungus with pores 23, 42
Fungus with spines 23
Calostoma 23, 29
Geaster 23, 28
Nidularia 23, 24
Clathrus cancellatus 24, 26, 121
Simblum rubescens 24, 26, 122
Anthurus borealis 24, 26, 121
Mutinus 27
Phallus impudicus 24, 27, 119

PAGE

Embryo plant of Mutinus caninus 24, 120
Scleroderma vulgare 25
Section of Scleroderma vulgare 25, 134
Mutinus caninus, young plant; embryo plant . . . 120
Bovista 28, 128
Section of Bovista (diagrammatic) 28, 128
Calvatia 28, 126
Section of Calvatia 28, 126
Geaster 28
Lycoperdon 29
Section of Lycoperdon (diagrammatic) . . . 29, 124
Bovistella 29
Section of Bovistella 29, 129
Clavaria 30
Section of Hydnum 30
Cap of Hydnum imbricatum 30, 96
Cap with striations on the margin 30, 40, 50
Bracket fungus 31
Resupinate fungus 31
Boletus Satanus 31
Polyporus quercina 31
Pleurotus, stem central 32
Pleurotus, stem eccentric 32
Pleurotus, stem wanting 32
Gills toothed, Lentinus 32
Lamellæ entire 33
Stem eccentric 33
Lamellæ simple 33
Fungus with volva and annulus 34
Volva, but no annulus 34
Annulus, but no volva 34
Lamellæ free from stem 34, 39
Annulus movable 34

List of Illustrations in Text

	Page
Lamellæ united with stem	34
Lamellæ decurrent	35, 37, 39, 40
Lamellæ adnate	35, 37, 39
Lamellæ sinuate	35
Volva wanting; annulus wanting	35
Lamellæ in shallow folds	36
Volva present; annulus wanting	37
Volva none; annulus none.	37
Annulus arachnoid	38
Lamellæ adnate	38
Veil remaining on stem as annulus	39
Volva none	39
Veil attached to margin of pileus	39
Lamellæ deliquescent	40
Spores fusiform	40
Spores globose	40
Hydnum with central stem	41
Branched Hydnum	41
Lamellæ labyrinthine, woody	43
Lamellæ radial, woody	43
Pores long-hexagonal	43
Pores in the form of tubes, stem lateral	44
Pores in the form of tubes, stem central	44
Section of young Amanita in wrapper	47
Young Amanita in wrapper	47
Young Amanita in ruptured wrapper	47
Section of Amanita muscaria	47
Cap striate, and free from warts	50
Section of Amanita Cæsarea	50
Young plant of Amanita Cæsarea	50
Section of Cantharellus floccosus	53
Section of Amanitopsis vaginata	54
Section of Amanitopsis parcivolvata	55

List of Illustrations in Text

	Page
Section of Lentinus lepideus	56
Section of Pleurotus ostreatus	57
Section of Pleurotus sapidus	58
Section of Pleurotus ulmarius	59
Section of Hygrophorus miniatus	60
Section of Hygrophorus eburneus	61
Section of Armillaria mellea	62
Section of Lepiota procera	63
Section of Lepiota naucinoides	64
Section of Collybia familia	67
Section of Russula emetica	68
Section of Russula virescens	69
Section of Clitocybe laccata	71
Section of Tricholoma personatum (var. bulbosum)	72
Section of Craterellus cantharellus	73
Section of Agaricus Rodmani	76
Section of Agaricus abruptus	77
Section of Hypholoma perplexum	79
Section of Hypholoma incertum	80
Section of Hypholoma sublateritium	81
Section of Pholiota adiposa	83
Section of Pholiota præcox	84
Section of Cortinarius violaceus	86
Section of Pluteus cervinus	87
Section of Lactarius piperatus	93
Section of Hydnum repandum	95
Section of Hydnum caput-ursi	96
Boletus Satanus	107
Under surface of Lenzites betulina	113
Favolus areolarius	115
Section of young phallus	119
Mutinus bambusinus	120
Lycoperdon	124

List of Illustrations in Text

	Page
Section of Lycoperdon (diagrammatic)	124
Asci and paraphyses	135
Section of truffle to show position of asci	135
Truffle, ascoma	135
Fungus growing on caterpillar	136
Section of Peziza odorata to show two layers	137
Gyromitra esculenta	141
Morchella esculenta	141
Section of Morchella deliciosa	141
Helvella	142
Helvella lacunosa (diagrammatic)	143

THE MUSHROOM BOOK

CHAPTER I: THE HOMES AND HABITS OF FUNGI

For centuries epicures have used certain fungi for food. The Greeks and Romans esteemed them highly, and gave a great deal of consideration to favourable times and places for gathering them, and to choice methods of preparing them for the table. Juvenal tells us of one old Roman enthusiast who was so carried away by his love for them as to exclaim, "Keep your corn, O Libya, unyoke your oxen, provided only you send us mushrooms!" Horace says that mushrooms which grow in the fields are the best, and that one can have but little faith in other kinds. Mushroom eaters of the present day would perhaps not agree with him, for they find edible species in every imaginable place where fungi grow, and are constantly adding to their list new varieties which they esteem delicious.

Although for centuries it has been known that some fungi contain most virulent poisons, still, through ignorance of those points which distinguish the poisonous from the edible, frequent cases of poisoning occur in all classes of society. The mistakes resulting in death have been frequent enough to inspire the timid with an overpowering dread of all fungi, while the damp and grewsome places in which many fungi flourish have caused them to be despised by others. The following lines from Shelley very aptly express the general sentiment:

> "And plants, at whose names the verse feels loath,
> Fill'd the place with a monstrous undergrowth,
> Prickly and pulpous, and blistering and blue,
> Livid, and starr'd with a lurid dew.

" And agarics and fungi, with mildew and mould,
Started like mist from the wet ground cold ;
Pale, fleshy, as if the decaying dead
With a spirit of growth had been animated."

SHELLEY: " The Sensitive Plant."

To many people the only growths known as fungi are toadstools and mushrooms. They give the name mushrooms to the species known to them as edible, and regard all other similar growths as toadstools, things uncanny or poisonous.

" The grisly todestool grown there mought I see,
And loathed paddocks [toads] lording on the same."

SPENSER'S " Faerie Queene."

This distinction has no scientific basis, and in fact most of the species called toadstools are edible. Fungi are not always the grewsome things of Shelley and Spenser. In their ranks are many which delight the eye with their colouring and the symmetry of their forms. They are the grotesques of nature; nests, hoofs, cups, umbrellas, shells, and clubs are represented, together with spheres, hemispheres, cones, and many other geometrical figures. The mildew on the linen, the mould on food, the rusts and smuts which blight our fields of grain, and the dry rot which crumbles our lumber to dust and which causes old wood in dark places to glow with a weird, pale, flickering light, are all forms of one group or another of these plants which prey upon living or dead organic matter. In ordinary observation, only the simpler and more noticeable fungi are taken into account, but they are in reality met with in almost every situation imaginable. They are found in damp cellars and in rooms shut off from the light ; in fact, some form of fungus will be found in every place and on everything which is not exposed to a circulation of fresh air.

In woods and open fields the attractive forms are found. In shady woods the beautiful white "bear's head" hangs on stately tree trunks, and the "destroying angels" gleam white in the shadows on the ground. Shelving brackets, green or red or brown, encircle old stumps, or stand out stiff and white from the crumbling trunks of fallen moss-grown monarchs of the forest, while wood-brown toadstools huddle in groups among

the fallen leaves. On the outskirts of the wood, green and red Russulas vie with the flowers in the brilliancy of their colouring. Pink or violet Clavarias, dainty corals, border the wood path, and golden Clavarias lighten up the sombre wood tints with their yellow branches. In dry pastures and along wood roads, puff-balls, large and small, send up their puffs of brown smoke, to the delight of every passing child who strikes them with a wand. On lawns and hillsides the Oreades cause fairy rings to grow. The fairy rings are circles, or parts of circles, of impoverished grass of a lighter colour and less luxuriant growth than that of the grass immediately surrounding the circle. Before the existence of fairy folk came to be doubted, it was firmly believed that these fairy rings were the dancing grounds of the fairies.

> "The nimble elves
> That do by moonshine green sour ringlets make
> Whereof the ewe bites not; whose pastime 'tis
> To make these midnight mushrooms."
>
> Rev. GERARD SMITH.

The rings on the commons increase in size until sometimes two or more rings intersect to form a labyrinth of green network. Rings appear year after year in the same place, and then disappear, to reappear after an interval of a few seasons. As long as the fairies existed in the imaginations of the people, it was easy to account for these strange happenings—the fairies danced in the moonshine, and the grass was worn down under their feet. If they were displeased and left the neighbourhood, the rings disappeared too. As this fancy was given up, other solutions of the mystery were sought. Some believed that the ring was caused by a thunder-bolt entering the ground at this spot, and still others were confident that it was caused by moles. The true solution is not hard to find, to one familiar with the habit of growth of the fungus plant. One fungus plant growing alone upon the lawn will soon exhaust the soil directly beneath it of all true fungus food. Of all the spores which fall from the parent plant only those will grow which fall without this impoverished spot, and so a ring of toadstools is formed. Again, only those spores which fall outside the ring will find good fungus food, and so the ring widens always outward, forming a perfect circle, unless something on one side or other interferes with its travels. The

decaying ring of fungi temporarily stimulates the grass around it, so that its rich colour stands out in circles or arcs of circles against the less highly nourished grass. Such rings are conspicuous on the lawns of the White House at Washington, and are often to be seen well defined on distant hillsides.

Brackets and mushrooms and puffballs grow in warm, moist places where they find decaying wood and leaves to feed upon. Old tree trunks and fallen logs, rich leaf mould, and cattle pastures are their favourite haunts.

The reason for their choice of place is invariably connected with the question of food, for fungi can thrive only where they can obtain organic matter, as they have lost the power which all green plants have of feeding on inorganic or mineral matter. All plants must have food with which to form plant flesh. Green plants by means of their leaf green—the only agent in the world which has the power to turn lifeless mineral matter into living matter—take the element carbon from the air, and hydrogen gas and oxygen gas from water, and with their green granules, by some mysterious process, make of the elements hydrogen, oxygen, and carbon, compounds of wood and starch and sugar. Fungus plants have none of this leaf green and must therefore feed on material which has been manufactured by green plants.

To define fungi simply, so as to include all the varieties, would be a difficult task ; but in general it may be said that they are plants which have no leaf green and which do not grow from true seeds, but from dustlike bodies resembling in appearance the yellow pollen of roses or lilies.




Puffball

The fungi have no flowers and produce no seeds. They produce spores instead, fine dust-like particles, which are borne in special places on the mature plant, whether a mould or mildew, a toadstool, puffball, or bracket. The cap of a mushroom placed right side up on a piece of paper under an inverted glass will print with its spores a picture of the radiating leaves or gills beneath. A slight blow on a puffball in the pasture will cause a puff of smoke-like dust to rise from it—really millions of spores that have ripened inside the puffball and are now ready to grow into new puffball plants when they fall on favourable soil.


CHAPTER II: THE RELATION OF FUNGI TO OTHER PLANTS

A CLASSIFICATION or orderly arrangement of material collected for study is indispensable to true pleasure and profit. The nature student must classify both his specimens and the knowledge he may obtain about them; for, as Spencer has said, "When a man's knowledge is not in order, the more of it he has the greater will be his confusion of thought." As he compares his specimens he sees interesting gradations of resemblance, and becomes fascinated with the pleasure of tracing their relationships and the gradual evolution of higher forms from lower.




Corollas and honey, attractive to insects

Every lover of nature who haunts the fields and woods acquires a rich store of facts about plant life, and without, perhaps, recognising that he does so, distinguishes two great groups of plants—those which have attractive flowers, and those which have no flowers at all. His flowerless plants bear no seeds, but quantities of fine, dust-like particles which rise in the air as he brushes his stick over their green leaves. As the powers of observation develop, he distinguishes the ferns and Christmas greens among flowerless plants, and perhaps soon recognises that the soft green moss bank, too, is composed of small plants, and that the green mats, the liverworts, on stones and moist banks and logs, are plants also. His only reason, perhaps, for calling them plants is that they grow and are green. He may



Ingenious stamens

learn with the microscope that the pond scums which he had thought disgusting frog-spittle are in truth tangles of exquisite plants, made up of chains of slender, transparent cells finer than silken threads, each cell containing many tiny green particles of leaf green, or chlorophyll—the cause of the green colour of all green plants.



Pistil of violet — Pistil of St. Johnswort

At first the most conspicuous plants attract the attention, and afterwards, in succession, those less and less conspicuous. They, in reality, present themselves in great natural groups, readily distinguished by well-marked characteristics.



Seed-box of iris



Seed-box of sacred bean

It will be seen, as these pass in review, that they are conspicuous according as they are complex. The gorgeous flowering plants have complicated methods of reproduction—corollas and honey, attractive to insects; ingenious stamens, pistils, seed-boxes, and seeds. The humble grasses, with their close relatives, dispense with gay colours and the assistance of insects, and trust to the breezes to carry their pollen to its goal. The pines and their allies are a step nearer simplicity, and do not enclose their seeds in a seed-box at all, but provide them with wings for dissemination, and leave them exposed to the wind.



Winged seed of the silver fir




Fern with spores (*Polypodium vulgare*)

The ferns and Christmas greens (Lycopodiums) have no flowers, and therefore no true seeds. They have a distinct stem, which grows from the apex and is strengthened by woody fibres, which may readily be seen by breaking the stem across. The woody fibres so strengthen the tissues of these plants that they are able to stand erect and make a conspicuous appearance not possible to the small mosses and liverworts, which are spore-bearing plants with no woody fibre.



Liverwort (*Marchantia polymorpha*)




Grass spikelet

The plants of all these groups resemble each other in descending degrees, so that they may be classed in groups under groups. Similar specimens may form groups of species. Species may form larger groups, or genera. Genera with common characteristics may form families ; and groups of families, orders; and orders, classes; while classes unite to form branches, or phylæ.



Indian pipe

A botanist relies for the classification of his specimens mainly upon the similarity of those parts of the plant which produce the seeds or spores rather than upon those parts—the roots and leaves and stems—which have the work of the plant household to do. He finds that the seed and spore producing parts are more constant in their forms and habits than the leaves and stems and roots, which are more exposed, and which are constantly



Bean seed open to show embryo

being forced to a change of form which will better suit their changed surroundings.

The novice sees nothing in the brown, or even in the highly coloured, fungi to warrant his calling them plants. They are to him "just toadstools;" for green colouring matter—his first criterion for plants—is not there, and, moreover, there is nothing in their shape which suggests to him the plants with which he is familiar. The snow-white Indian pipe lacks the green of most plants, but that does not rule it for him out of the plant world; for although it is colourless, and depends upon other plants for food, still it has a flower form and produces a seed-box with well-developed seeds. Fungi, however, to any but the close student must seem quite unrelated to all normal plant forms. But the botanist, by a study of their structure, finds that they all grow from microscopic, dust-like particles, which differ from true seeds in consisting of but one or a few cells, and in having no embryo plant in them as true seeds have. He recognises their position in the kingdom of living things, and classes them as spore-bearing plants, lower than the group of mosses, those dainty plants which delight every one with their gracefulness, and which bear their spores in tiny capsules or boxes set up on slender stems. By studying their life history he decides that they are degenerate members of the lowest group—the algæ—and that they have fostered the habit of feeding on material constructed by green plants, instead of constructing food material for themselves, and have, in consequence,



Moss (*Dicranum scoparium*) (natural size)




White mould on dead fly

lost their power of constructing such food, and also their green granules by which this work of construction may be carried on. The life history and structure of fungi has been studied so minutely that one is able to arrange them in three well-marked classes:




Bread mould (magnified)

The first class, the algal-like fungi (Phycomycetes), includes bread moulds and several of those fungi which cause diseases of plants and animals—the downy mildew on the grape, the potato rot, the common white mould which fastens dead flies to the walls or window panes in the autumn, and the fungus which grows on salmon and causes them to die in great numbers. The plant of these fungi is cobwebby, sometimes growing within the cells of the plant substance on which it lives, and sometimes growing both within and on the surface. A freshly moulded piece of moist bread shows the bread covered with exquisitely fine transparent threads, which constitute the plant. Later, spore cases containing tiny black spores will be seen, which give a delicate gray tint to the plant at first, but later form a black, repulsive mass as their numbers increase. These plants are regarded as descendants of degenerate algæ, which lost their power of independent existence through stealing their food instead of making it for themselves.


Spores borne in delicate membranous sacs (magnified)

The second class, the spore-sac fungi, produce their spores in delicate membranous sacs. The spore-sac fungi vary greatly in size, habit, and structure. Most of them are inconspicuous members of the plant world, as the yeast plant, by which our bread is raised; the fungus which causes the peach leaves to curl and the black knots to appear on cherry and plum trees.


Spores borne on little spicules (magnified)

The third class is made up of all fungi which bear their

spores on little spicules standing up on large cells. This contains most of the conspicuous fungi one will care about knowing. To understand the group one must understand the method by which a spore grows to be a fungus plant, and to be able to distinguish the different members of the group one must know on just what portions of the spore receptacle the spores are borne.

CHAPTER III: FROM SPORE TO MUSHROOM

THE way in which a spore grows into a fungus plant is very simple:

(1) The spore is a single cell, and when it is in a warm, moist place it swells.


(2) The cell absorbs food through its cell wall and divides into two cells.



(3) Each new cell absorbs food and divides until long chains of cells are formed, looking to the unaided eye like threads. Each thread is a hypha, and a tangle of threads is a mycelium.



(4) In the soil the mycelium nourishes itself on decaying vegetable matter, and grows; then, at certain points, the threads mat together to form little balls the size of pinheads (*a*).




(*b*) The pinheads grow to the size of bird-shot.



(*c*) The bird-shot increase to the size of shoe-buttons.

(5) If the ball is to become a stemmed toadstool, a minute stem appears on the button. The stem and button increase in size. The button is lifted above the soil and expands into a mushroom.




(6) If the button is to become a puff-ball, no stem appears on the button; but

it grows, and comes out of the ground a round puffball. (See Plate opposite p. 124.)

If one wishes to learn to distinguish the members of the mushroom or toadstool family, either for the pleasure he may derive from knowing them, or from a desire to distinguish the edible from the poisonous, he must be familiar with the typical parts of the fungus plant, and must know the names of these parts.



Puffball



The edible mushroom of the market (*Agaricus campestris*) serves well for study, as it shows some of the characteristics which all the toadstools, mushrooms, brackets, and puffballs have in common.



(1) This mushroom is in shape something like a parasol.

(2) The handle is the stem, or stipe.

(3) The open top is the cap, or pileus.



(4) Under the cap, radiating from the stalk to the edge of the cap, are thin plates—the gills, or lamellæ.




Section across gill (magnified)

(5) When the mushroom is in the button stage, the gills are not visible, for they are covered with a thin sheet of mycelial threads, called the veil. (See coloured plate of *Agaricus campestris*.)

(6) As the button grows the veil stretches, and finally breaks,

leaving a ragged edge to the cap, and a ring or annulus of veil around the stem. The gills of the *Agaricus* are not fastened to the stem, but are rounded off at the end near the stem, while others, between the long ones, extend from the edge of the cap only far enough toward the stem to fill up the angles formed by the long gills.



A small portion of section of gill (highly magnified)

The surface of the gills is the fruiting portion of the mushroom. It is here that the spores are formed.

The structure of the fungus plant up to this point has been similar throughout. A loose tangle of threads underground formed the mycelium—the food provider. A more closely matted tangle above ground formed the stem and cap and veil, and even the central part of the gill—the fruiting parts of the plant.



Corn smut

On the surface of the gill a difference in structure is found, which will be clearly understood from a picture of a thin section cut across a gill.

(1) The central portion of the gill is made by loosely tangled mycelium threads (*tr*) draping themselves in thin plates from the surface of the cap.


(2) Just outside of this loose mycelium, on either side, are layers of short cells (*c*), which bear club-shaped bodies standing out over both surfaces of the gills (*b*).

(3) Each club bears two slender processes (*st*) at the free end, and each process bears a spore (*sp*).



Leaf rust on *Hepatica triloba*

All corn smuts, wheat smuts, leaf rusts, toadstools, puffballs, and brackets bear their spores on club-like cells, and for this reason are put in one group, called Basidiomycetes.




Pouch-fungus section, to show spores in hollow rind

The fact that corn smuts and leaf rusts feed on living plants, while toadstools, brackets, and puffballs feed on dead plants, separates them into two groups; the smuts and rusts forming the lower group, and the others the higher group. It is the higher Basidiomycetes which we wish to consider, as this group includes most of the conspicuous fungi, most of the edible, and those fungi which are dangerous because of their resemblance to edible species.



Section to show gills




Section of a Boletus, to show pores

Remembering that toadstools, puffballs, and brackets all start from spores; that all have the tangled thread-like plants, seeking the dark; that they all have the spore receptacle in the light, and bear their spores on club-like cells, one can readily understand their being put in one group.



Clavaria with spores on spines



Section of Hydnum, to show teeth

With a few exceptions not necessary for us to consider, all the higher fungi naturally divide into two groups—pouch-fungi (Gasteromycetes), which conceal their spores in a definite rind, or peridium, as the puffballs do; and membrane fungi (Hymenomycetes), now called Agaricales, which bear their spores exposed on the surface of gills, pores, spines, or teeth, as the garden mushrooms, the Boleti, the Clavarias, and the Hydnums.

CHAPTER IV: THE KEY

WHAT A KEY IS, AND WHY A NAME IS DESIRABLE

A KEY in the study of botany is a guide by which a student may trace a specimen until he finds a name for it. Having found a name, he may learn from books or from friends what is known of its habits of growth, of its value as a food or drug, whether it is harmful or harmless, whether it is to be protected or whether war is to be waged against it. He may learn whether it has figured in history or the myths, and how the poets and artists viewed it, and may perhaps learn to see it with their eyes. He may watch similar specimens as they grow, and may add the results of his observations to the facts already recorded about his specimen.

HOW A KEY FOR FUNGI IS MADE, AND WHY IT IS DESIRABLE

In the first place, only such plants are considered as grow from spores and have no leaf-green. (The spore characteristic is one the amateur must decide upon either by seeing the spores or by inferring their existence from the fact that seeds do not appear.) There are some thirty-five thousand species of fungi known to botanists, so that it would be impossible to find a name for a specimen if one had to read at random until the right description for his specimen was found; but since all of these plants may be put in one or another of three groups, on account of certain points of resemblance which they have in common, and since these three groups may each in turn be divided and subdivided, one may, by selecting groups rather than individual specimens, find a short path to the name desired. The three primary groups, called *classes*, are made as follows :

The first contains many mould-like fungi which resemble one another in microscopic characters.

The second contains other mould-like fungi and many con-



spicuous fungi which bear their spores in transparent sacs (see first page of Key).

The third contains all fungi which bear their spores on enlarged cells called basidia (see first page of Key).

To even partially understand the inconspicuous fungi is a task impossible to one who is not familiar with the use of a compound microscope. To acquire a knowledge sufficiently accurate to identify nearly all of the conspicuous fungi is within the power of any intelligent person, for the two groups or classes containing the conspicuous species may be divided, on account of easily distinguished characters, into groups called *orders.* The orders may be divided into groups called *families,* and the families into groups called *genera* (singular genus), and the genera into individual specimens called *species;* and all these groups may be arranged in such a way that the series of selections may be quickly made. Such an arrangement of groups is called a key.

KEY



FUNGI.


		CLASS
Fungi which are inconspicuous as individuals, and do not bear the spores in sacs (asci), or on sterigmata. Mould-like.		I Phycomycetes. (The *Algal-like Fungi.*) Page 9.
Spores produced in delicate membranous sacs (asci).	 (Magnified)	II Ascomycetes. (The *Spore-sac Fungi.*) Page 18.
Spores free on enlarged basidia.	 (Magnified)	III Basidiomycetes. Page 22.

Singular, fungus ; pronounced fŭn′-gŭs. Plural, fungi ; pronounced fŭn′-jī
" ascus ; " ăs′-kŭs. " asci ; " ăs′-sī
Phȳ′-cō-mȳ-çē′-tēs Ăs′-cō-mȳ-çē′-tēs Bă-sĭd′-ĭ-ō-mȳ-çē′-tēs

CLASS II

ASCOMYCETES (SPORE-SAC FUNGI)

		ORDERS
+ Subterranean. Asci remaining enclosed in a tuber-like body.	 Outer surface  Section	Tuberales. (Truffles.) Page 135.
+ Bright coloured. Perithecia imbedded in a fleshy or waxy stroma. Asci collected in a spherical or pear-shaped body.	 Section of stroma with perithecia (magnified)  Perithecium (highly magnified)	Hypocreales. Page 136.


+ Fleshy cup-like fungi. Asci collected in a flattened or cup-like body (ascoma). Ascoma closed at first, open at maturity.		Pezizales (Cup Fungi.) Page 20.
+ Club-shaped (*a*); or conic, convex, and pitted (*b*); or with gyrose furrows (*c*); or saddle-shape (*d*).  (*c*)	 (*b*)  (*d*)  (*a*)	Helvellales. Page 20.

Singular, pĕr-ĭ-thē′-çĭ-ŭm ; plural, pĕr-ĭ-thē′-çĭ-ă
" ăs-cō′-mă " ăs-cō′-mă-tă

Tū-bĕr-ā′-lēs Hўp′-ŏ-crē-ā′-lēs Pēz′-ĭ-zā′-lēs Hĕl′-vĕl-lā′-lēs

CLASS II

ASCOMYCETES (SPORE-SAC FUNGI)

					GENERA
⊙ Pezizales.		♀ Ascomata smooth, regularly saucer-shaped, or cup-shaped, or circular.			Peziza. Page 137.
⊙ Helvellales.	× Family Helvellaceæ.	♀ Ascoma conic or pileate.	⊕ Ascoma with a distinct stalk, campanulate or saddle-shaped, attached to the stipe at the middle.		Helvella. Page 142.
			⊕ Ascoma cap-like, irregular or lobed, covered with gyrose wrinkles.		Gyromitra. Page 141.

Family Geoglossaceæ.

× ♀ Ascoma clavate or capitate; yellow, green, or black club-like forms.

⊕ Ascoma cap-like, ovoid, or conic, covered with deep pits.

Morchella. Page 141.

⊕ Ascomata flat, running down both sides of the stem.

Spathularia. Page 138.

⊕ Ascoma hollow, discoid, usually with free margin, light-coloured, yellowish or light brown, sharply separate from the stem.

Mitrula. Page 140.

Hĕl'-vĕl-lā'-çĕ-æ Pĕz-ĭ'-zā Hĕl-vĕl'-lā Ġў'-rō-mĭ'-trā Ĝĕ'-Ӑ-glŏs-sā'-çĕ-æ Mŏr-kĕl'-lā Spāth'-ū-lā'-rĭ-ā Mĭt'-rŭ-lā


CLASS III

BASIDIOMYCETES (SPORES BORNE ON BASIDIA)




▲ Fungi taking nourishment from living plants, parasitic, often deforming the host.	 			Rusts and Smuts. Page 14.
	Gelatinous fungi with divided spore clubs.			Tremellines. (Gelatinous Fungi.)
▲ Fungi taking nourishment from dead organic matter.	○ Fleshy, woody, or leathery. Spore clubs undivided.	+ Spore clubs forming a membranous surface, naked at maturity. 1st. Membrane covering the surface of lamellæ.		Hymenomycetes, or Membrane Fungi, or

2d. Membrane covering the surface of pores.




Agaricales. Page 30.


3d. Membrane covering the surface of spines.



+ Spore clubs enclosed within a definite case (peridium).









Gasteromycetes. (Pouch Fungi.) Page 24.




Trĕm′-ĕl-lĭnes Hȳ′-mĕn-ō′-mȳ-çē′-tēs Pĕr-ĭd′-ĭ-ŭm Ā-găr-ĭ-cā′-lēs Găs′-tĕr-ō′-mȳ-çē′-tēs


GASTEROMYCETES (POUCH FUNGI)		ORDERS
□ Spores borne in a more or less deliquescent mass, which is at first enclosed in an egg-like sac (peridium), but at maturity is elevated on an elastically expanding receptacle.		Phallales. (Stinkhorns.) Page 26.
□ Spores borne in little egg-like cases, persistent, in a cup-shaped peridium.		Nidulariales. (Bird's-nest Fungi.) Page 133.

◘ Spores remaining within the peridium until maturity.	◡ Fleshy when young; at maturity filled with dust-like spore masses, mixed with threads (capilitium).		Lycoperdales. Page 28.
	◡ Rough, warty balls, thick skinned. Lead colour within. Skin rupturing irregularly at maturity.	  Section	Sclerodermatales. Page 133.

Phăl-lā′-lēs Nĭd′-ū-lă′-rĭ-ā′-lēs Căp-ĭl-lĭ′-shĭ-ŭm Lȳ′-cō-pĕr-dā′-lēs Sclĕr′-ŏ-dĕr′-mă-tā′-lēs




ORDER. PHALLALES (STINKHORNS)

				GENERA
≡ Spores borne on a more or less deliquescent mass (gleba), which is at first enclosed in an egg-like sac (peridium), but at maturity is elevated on an elastically expanding receptacle.	‖ Receptacle latticed or irregularly branched, sessile or stalked; spore mass enclosed by the receptacle.	Clathraceæ.	= Receptacle sessile; latticed with columnar or obliquely anastomosing bars. 	Clathrus. Page 121.
			= Receptacle latticed; meshes of about equal diameter either way. 	Simblum. Page 122.
			= Receptacle ending in free arms which enclose the gleba when young, but afterwards diverge. 	Anthurus. Page 121.

Phallaceæ.	‖ Receptacle *not* latticed; spore mass *not* enclosed by the receptacle.	= Gleba borne on the upper portion of the stem.			Mutinus. Page 120.
		= Gleba borne on the surface of a pileus which joins the stipe at the apex.	∞ Veil hanging from the apex of the stipe underneath the pileus.		Dictyophora. Page 120.
			∞ Veil wanting.		Phallus. Page 117.

Glē′-bă Clăth-rā′-çē-æ Clăth′-rŭs Sĭm′-blŭm Ăn-thū′-rŭs Phăl-lā′-çē-æ Mū-tī′-nŭs Dĭc′-tĭ-ŏph′-ŏ-ră Phăl′-lŭs


Key

ORDER. LYCOPERDALES		GENERA
Ψ Peridium sessile. No sterile base. Peridium opening by an apical mouth.	 Section	Bovista. Page 128.
Ψ Peridium sessile, with sterile base. Upper surface breaking into fragments from above downwards.	 .....Sterile base Section	Calvatia. Page 126.
Ψ Peridium splitting into starlike segments. Peridium opening by a single mouth.		Geaster. Page 129.

Ψ Peridium with sterile base.


Peridium opening by an apical mouth. Capillitium (threads) present.

λ Threads long, slender, attached.




Lycoperdon. Page 124.

λ Threads short and free.





Bovistella. Page 129.

Ψ Outer peridium gelatinous, and later disappearing. Base of interlacing fibres. Red star-like mouth.



Calostoma. Page 131.

Bŏ-vĭs'-tă Căl-vā'-shĭ-ă Ġĕ-ăs'-tĕr Lȳ-cŏ-pĕr'-dŏn Bŏ-vĭs-tĕl'-lă Căl-ŏs'-tŏ-mă

ORDER. AGARICALES		FAMILIES
[1] Spores borne on upper portions.	φ Club - shaped, or forming masses of erect branches rising from a common base. 	Clavariaceæ. Page 98.
[1] Spore - bearing surface normally (1st) provided with cap and central stem. 	φ Spores borne on radiating plates or lamellæ. 	Agaricaceæ. Pages 32-40.
	φ Spores borne on teeth, tubercles, or tooth-like plates projecting downward in growth. 	Hydnaceæ. Page 41.

Section

(2d) Bracket-like.




(3d) Resupinate.



φ Spores borne on the interior of pores or tubes or labyrinthine passages.

* Substance fleshy. Pores more or less easily separating from the cap and each other.

Stem central, or stem lateral.



Boletaceæ.
Pages 44, 45.


* Substance corky, woody, or leathery.



Polyporaceæ.
Pages 42, 43.


Clăv'-ăr'-ĭ-ă'-çĕ-æ Ă-găr'-ĭ-cā'-çĕ-æ Hў̆d-nă'-çĕ-æ Bō-lĕ-tă'-çĕ-æ Pō'-lў̆-pō-rā'-çĕ-æ

FAMILY. AGARICACEÆ (FUNGI WITH GILLS)

			GENERA
	Juice milky, white or coloured.		Lactarius. Page 92.
⊥ Plant fleshy; soon putrescent.	± Juice watery.	+ Stem: Lateral, 1. Eccentric, 2. Wanting, 3.  1 2 3 Spores white. (Spores violet in P. sapidus.)	Pleurotus. Page 57.
		+ Stem central. Spores white	Pages 34, 35, 36, 46-73.
		Spores rosy or salmon	Pages 36, 37, 87, 88.
		Spores yellowish-brown or rusty-brown	Pages 38, 83-87.
		Spores dark brown or purplish-brown .	Pages 38, 39, 73-83.
		Spores black	Pages 40, 89-92.
⊥ Plant tough, leathery, or woody; reviving or persistent.	± Juice watery.	+ Lamellæ normally toothed on their edges.  Stem central. Stem eccentric. Stem wanting.	Lentinus. Page 56.

3	+ Lamellæ entire. 	● Stem central.	☾ Lamellæ simple; cap firm and dry.		Marasmius. Page 65.
			☾ Lamellæ simple; plant leathery.		Panus. Page 67.
33		● Stem lateral or wanting.	☾ Lamellæ deeply splitting; villous.		Schizophyllum. Page 67.
			☾ Lamellæ channelled or crisped.		Trogia. Page 67.

Lăc-tā′-rĭ-ŭs Plū-rō′-tŭs Lĕn-tĭ′-nŭs Mă′-răs′-mĭ-ŭs Pā′-nŭs Skĭz-ō-phȳ′-lŭm Trō′-ġĭ-ă



	FAMILY. AGARICACEÆ (FUNGI WITH GILLS) (WHITE-SPORED SERIES)	GENERA
○ Volva and annulus.		Amanita. Page 46.
○ Volva, but no annulus.		Amanitopsis. Page 53.
○ Volva wanting; annulus present.	☿ Lamellæ free from stem; annulus often movable; cap usually scaly.  	Lepiota. Page 63.
	☿ Lamellæ united with the stem; pileus usually smooth. 	Armillaria. Page 61.




o Volva wanting; annulus wanting.	☿ Lamellæ thin, their edges acute.	× Lamellæ decurrent on the stem. 		β Stem fleshy.	Clitocybe. Page 70.
				β Stem with cartilaginous rind.	Omphalia. Page 68.
		× Lamellæ adnate.	θ Stem with cartilaginous rind. 		Collybia. Page 66.
			θ Stem fleshy; cap often bright-coloured.	ν Plant rigid; lamellæ usually brittle.	Russula. Page 68.
				ν Lamellæ waxy.	Hygrophorus. Page 59.
		× Lamellæ sinuate. 	θ Stem fleshy.		Tricholoma. Page 72.
			θ Stem with cartilaginous rind; cap membranous, more or less less striate.		Mycena. Page 55.

Ă-măn′-I-tă Ă-măn′-I-tŏp′-sĭs Rŭs′-sŭ-lă Lĕp-i-ō′-tă Hў-grŏph′-ŏ-rŭs Ăr′-mĭl-lā′-rĭ-ă Trĭ′-chŏ-lō′-mă Clĭ-tō-çў′-bē Mў-cē′-nă Ŏm-phā′-lĭ-ă Cŏl-lў′-bĭ-ă



Key

FAMILY. AGARICACEÆ (FUNGI WITH GILLS) (WHITE-SPORED SERIES)		GENERA
○ Volva none; annulus none.	Lamellæ in the form of shallow folds, their edges obtuse.  Lamellæ decurrent; plant terrestrial.	Cantharellus. Page 53.
FAMILY. AGARICACEÆ (FUNGI WITH GILLS) (PINK-SPORED SERIES)		
○ Volva present; annulus wanting.		Volvaria. Page 88.
○ Volva wanting; annulus present.	(Annularia has not been found in America.)	Annularia.



○ Volva none; annulus none.	∠ Lamellæ free from stem (*a*). 	Pluteus. Page 87.
	Stem fleshy.	
	∠ Lamellæ adnate (*a*) or sinuate (*b*). 	Entoloma. Page 88.
	∠ Lamellæ decurrent on the stem. 	θ Stem fleshy. — Clitopilus. Page 88.
		θ Stem with cartilaginous rind. — Eccilia. Page 88.

Căn-thă-rĕl′-lŭs Vŏl-vā′-rĭ-ă Ăn′-nŭ-lā′-rĭ-ă Plū′-tĕ-ŭs Ĕn′-tŏ-lō′-mă Clĭ-tŏ-pī′-lŭs Ĕc-çĭl′-ĭ-ă

FAMILY. AGARICACEÆ (FUNGI WITH GILLS) (RUSTY-SPORED SERIES)			GENERA
○ Annulus not arachnoid; continuous and persistent.			Pholiota. Page 83.
○ Annulus arachnoid, filamentous, or evanescent; often not apparent in older specimens.		Lamellæ adnate; plant terrestrial. 	Cortinarius. Page 85.

FAMILY. AGARICACEÆ (FUNGI WITH GILLS) (BROWN-SPORED SERIES)		
○ Volva at the base of the stem.	(Single species reported from Nebraska.)	Chitonia. Page 83.

o Volva none.

ff




ſ

μ Veil remaining on the stem as an annulus.



ρ Lamellæ free from the stem.



Agaricus. Page 73.

ρ Lamellæ united with the stem.



Stropharia. Page 82.

μ Veil remaining attached to the margin of the pileus; often not apparent in very old specimens.



Hypholoma. Page 78.

μ Veil inconspicuous or wanting.

ρ Lamellæ free.



Pilosace. Page 82.

ρ Lamellæ decurrent.



Deconica. Page 82.

ρ Lamellæ adnate.







Psilocybe Page 82.

Psathyra. Page 82.


Phŏ-lĭ-ō′-tă Cŏr′-tĭ-nā′-rĭ-ŭs Chĭ-tō′-nĭ-ă Ă-găr′-ĭ-cŭs Strŏph-ā′-rĭ-ă Hȳ-phŏ-lō′-mă
Pĭ-lŏ-sā′-çē Dē-cŏn′-ĭ-că Psĭ-lō-cȳ′-bē Psă-thȳ′-ră



Key



FAMILY. AGARICACEÆ (FUNGI WITH GILLS) (BLACK-SPORED SERIES)			GENERA
○ Lamellæ deliquescent, melting to an inky fluid.			Coprinus. Page 89.
○ Lamellæ not deliquescent; annulus wanting.	∠ Lamellæ waxy, decurrent. Spores fusiform.	 	Gomphidius. Page 92.
	∠ Lamellæ not waxy, not decurrent. Spores globose-ovoid.	 ω Pileus striate, with tough rind. 	Psathyrella. Page 92.
		ω Pileus not striate; stem fleshy.	Panæolus. Page 92.

41	2 Plant consisting of teeth attached to a resupinate membrane or to a pileus.	μ Species with central stem.				Hydnum. Page 95.
	2 Spore surface covering needle-like spines, awl-shaped, usually regular.	μ Species growing in branched masses, with no distinct pileus.				
	Cŏ-prĭ′-nŭs	Gŏm-phĭd′-ĭ-ŭs	Psă-thў-rĕl′-lă	Păn-æ′-ŏ-lŭs	Hўd-nă′-cĕ-æ	Hўd′-nŭm

Key

FAMILY. POLYPORACEÆ (FUNGI WITH PORES)				GENERA
✕ Spore-bearing surface not waxy; not separable from the context.	△ Pores minute and round, or larger and angular.	Σ Substance of the cap (trama) descending between the pores.	(This genus is with difficulty separated from the next).	Trametes. Page 113.
		Σ Substance of the cap not descending, namely, substance between the pores different from that of the cap.		Polyporus. Page 110.

▵ Pores large, long-hexagonal, sinuous, or lamelliform.	Σ Stem lateral; pores long-hexagonal.		Favolus. Page 115.
	Σ Lamellæ sinuous and labyrinthine; stem normally wanting.		Dædalea. Page 114.
	Σ Lamellæ radial; stem wanting, or spore-bearing surface growing flat on its host.		Lenzites. Page 113.
Tră-mē'-tēs	Pō-lўp'-ŏ-rŭs	Fā'-vŏ-lŭs	Dæ-dā'-lē-ă	Lĕn-zī'-tēs

	FAMILY. BOLETACEÆ (FUNGI WITH PORES)		GENERA
⁂ Pores quite easily separable from the pileus and each other.	⏢ Stem lateral; pores in the form of tubes, whose mouths are separated from each other.		Fistulina. Page 102
	⏢ Stem central, or, rarely, somewhat eccentric.	^ Pores readily separating from the pileus; spores brown or whitish. 	Boletus. Page 104.

	^ Pores separating with difficulty from the pileus.	⩛ Fleshy pores, in more or less radiating rows; spores brown or yellowish.	Boletinus. Page 103.
		⩛ Tough; pores uniform; pileus floccose; spores blackish.	Strobilomyces.
Fĭs-tŭ-lĭ′-nă	Bŏ-lē′-tŭs	Bŏ′-lĕ-tī′-nŭs	Strŏ′-bĭl-ŏ′-mў̆-çēs

CHAPTER V: FUNGI WITH GILLS

GENUS AMANITA

THE fungi with gills all have this characteristic in common—that they bear their spores on radiating plates or lamellæ. Their family name, *Agaricaceæ*, is derived from a typical member of the family, *Agaricus campestris*. The family is the largest and most widely distributed of all the families, and contains some five thousand described species, which are placed in groups or genera, more or less large, based on such characters as the colour of the spores; the position and shape of the lamellæ; the colour and texture, as well as the shape, odour, taste, and appendages of the cap and stem.

The colour of the spores is one of the most important characteristics, as the decision as to whether the plant is wholesome or not often rests upon it. The colour may be determined by placing the cap, with spore surface down, on a sheet of white or black paper, and leaving it for a time under an inverted glass, so as to cut off all drafts which may blow the spores away. A print of the radiating gills will then be made in the colour of the spores—white, pink, rusty brown, or black.

For external characters of the stem, one must be careful to get the entire stem from the ground; for a most important characteristic, the volva, if present, will be found at the base. The volva may be membranous and attached to the base, excepting at the rim, or membranous and loose, or present only in the form of rings of scales at the base, with perhaps traces on the surface of the cap.


The surface of the stem may be smooth or rusty or mealy white. It may or may not have near the cap a ring of the membrane which covered the gills of the young plant.

The character of the lamellæ and the internal characters of the stem may best be determined by cutting the cap and stem from top to base with a sharp knife, for then it may be seen whether the gills are free from the stem or attached to it, or

whether they grow down on the stem; and whether the stem is hollow, solid, or filled with web-like mycelium, and whether it is fleshy or has a tough and hard rind. The lamellæ may be of different or of equal lengths, and their edges may be entire or toothed or thin or blunt.


To know a genus of the gill-bearing fungi, one must know the cap, gills, stem, and habit of growth which characterise that genus.

A very young plant of the genus *Amanita* is enveloped in a membranous wrapper. The relation of the young plant to the wrapper will readily be understood by cutting a young plant through its length.



Young plant in wrapper


As the plant grows, the wrapper is ruptured, a part is left at the base to form a cup or sheath, or a part may be carried up on the cap, to appear in small patches.




Section of young plant in wrapper

The cap is, as a rule, regular and broadly convex. It may be almost flat when mature. The stem has a conspicuous collar and the gills are free from the stem.

There are twenty American species in the genus *Amanita;* some of them are the most poisonous fungi known, while others are most highly esteemed for the table. Since the most dangerous species belong to this genus, it would be better for the amateur not to eat of specimens which have stalks with a swollen base surrounded by a cup-like or scaly envelope, especially if the gills are white. In gathering all white-gilled species, care should be taken to get



A. muscaria (See Plate III.)



Wrapper ruptured

below the base of the stalk; for it often happens that the bulb is broken off and left behind, and thus the principal characteristic lost which would mark it as a specimen not to be eaten.

Death Cup; Poison Amanita (Poisonous)

Amanita phalloides

Cap or Pileus—White or greenish or greyish brown; smooth, no striations; width, 3–5 inches.
Stem or Stipe—Ring present. Abruptly bulbous at the base; bulb margined by the wrapper remains. White in white-cap forms, tinged with a paler shade than the cap in brown-cap forms. Pithy when young, hollow when old. 3–6 inches long.
Veil—White in white-cap forms, tinged with brown in brown-cap forms.
Gills or Lamellæ—White, free from the stem, rounded at the stem end, rather broad.
Spores—Globose and white.
Flesh—White.
Time—July to October.
Habitat—Woods, groves, open places, and pastures.

The poisonous principle of the death cup is known as phallin, one of the tox-albumins, the poisons found in rattlesnakes and other venomous animals, and the poisons which produce death in cholera and diphtheria.

The phallin acts directly upon the blood corpuscles, dissolving these, so that the serum of the blood escapes from the blood-vessels into the alimentary canal and drains the whole system of its vitality. There is no known antidote by which the effects of phallin may be counteracted. If one has eaten of the *Amanita phalloides*, the only chance of saving life is to remove the undigested parts from the alimentary canal by stomach-pump and oil purgatives; then, if the amount of phallin absorbed into the system is not too great, the remainder may wear itself out on the blood and the patient may recover.

The amount of the fungus which is necessary to produce death is small; even the handling of specimens and the breathing in of spores affect some people unpleasantly.

Phăl-loi'-dēs



DEATH CUP. DESTROYING ANGEL

(*Amanita phalloides*, Fries)

Reduced. Nat. size: Cap, 3½ inches; stem, 7½ inches



Fly Amanita (poisonous).
(Amanita muscaria, L.).

Fly Amanita (Poisonous)

Amanita muscaria

Cap or Pileus—Orange red to pale yellow or almost white. The young plants are brighter, and fade from the margin inward as the plant matures. Floccose scales, the wrapper remains, are scattered on the cap. The margin is often striate. 3–6 inches broad.

Stem or Stipe—White or slightly tinged with yellow. Pithy or hollow. Base not broad and abrupt, but ovate, covered with the scaly margins of the wrapper. 4–6 inches long.

Veil and Ring or Annulus—The veil covers the gills of the young plant, and later is seen as a collar-like ring on the stem.

Gills or Lamellæ—White or slightly tinged with yellow. Various in length ; short ones terminating in length with almost vertical abruptness.

Spores—White, broadly elliptical.

Flesh—White, tinged with yellow under the epidermis.

Habitat—Along roadsides, on borders of fields, in groves of coniferous trees. It prefers poor soil, gravelly or scanty. It grows singly, not in groups.

Time—June until freezing weather.

Young Plant—This is at first egg-like, then dumb-bell shaped. As the parts within expand, the wrapper breaks up into scales, so that the convex, unexpanded cap is densely covered with more or less concentric fragments of the wrapper, and the bulbous stem is covered with rings of fringy scales. As the stem expands, these scales are left on the bulbous base, while the fragments on the cap are more widely separated by the growth of the cap.

The fly amanita is a very conspicuous and handsome species. There are conflicting statements concerning the properties of this fungus; some claim that it is edible, and yet it is known to have caused much sickness and many deaths. It caused the death of the Czar Alexis of Russia, and of the Count de Vecchi in Washington. It is said that it is cooked and eaten by the Russians, and still it is on record that several French soldiers ate of it in Russia and became very ill.

The Siberians steep dried specimens of the fly amanita in whortleberry juice, and thus make a drink which produces an intoxication similar to that produced by the "haschisch" and "majoon" of the East.

Mŭs-că'-rĭ-ă

There is something about it particularly attractive to flies, and yet for them to sip its juices means death, as may be seen by the circle of dead flies lying on the ground under the shadow of its cap.

The chief poison of this fungus is an alkaloid called muscarine, which paralyzes the nerves controlling the action of the heart. Injections of atropine in doses of from one one-hundredth to one-fiftieth of a grain are employed as an antidote for this poison. In addition, the most powerful emetics are used.

Orange Amanita (Edible)

Amanita Cæsarea

Cap or Pileus—Smooth, glabrous, and free from warts or scales. Red or orange, fading to yellow on the margin or all over the cap. Margin distinctly striate. When fully expanded, nearly flat. When moist, slightly sticky and viscid.




Cap *striate*, and free from warts

Stem—Yellow. When young, fibrous or cottony within, later hollow.

Wrapper or Volva—White and membranous, loosely sheathing the base of the stem.


Veil—Covers the gills of the young plant. Remains are seen on the stem only, where it hangs down like a white ruffle.

Gills or Lamellæ—Rounded at the stem end and not attached to the stem. Yellow, an exception to the rule that the colour of the gills in mature plants resembles the colour of the spores.



Section of A. Cæsarea

(*a*) Smooth cap (*c*) Hollow stem
(*b*) Free gills (*r*) Ruffle-like ring



Young plant

Young Plant—When young, the cap and stem are contained in a wrapper not unlike a hen's egg in shape, size, and colour. As the cap and stem within develop, the wrapper ruptures in its upper part, the stem elongates, and the cap is carried up,





ORANGE AMANITA (EDIBLE)

(*Amanita Cæsarea*, Scop.)

Cap diam., 3-8 inches; stem length, 3-8 inches

while the remains of the wrapper are left at the base of the stem, an open sac.

Spores—White, elliptical.

Flesh—White stained with yellow under the separable epidermis and next the line of attachment of the gills.

Taste—Mild and pleasant.

Habitat—Thin woods, preferably pine woods and sandy soil. Abundant in southern Europe, common in the Southern States, and occasionally found in New York and Massachusetts.

Time—July, August, September.

The *Amanita Cæsarea* is one of the handsomest species. The Greeks and Romans esteemed it as an article of food. The names, "Food of the gods," "*Cibus Deorum*," "Imperial mushroom," "Cæsar's mushroom," and "*Kaiserling*," suggest the esteem in which it was held.

COMPARISON OF THREE YELLOW AMANITAS

Amanita Cæsarea (Edible)	Amanita muscaria (Poisonous)	Amanita Frostiana (Poisonous)
Cap or Pileus—Bright orange or, rarely, paler yellow. Smooth.	*Cap or Pileus*—Orange or yellow, adorned with flocculent warts consisting of patches of the ruptured volva.	*Cap or Pileus*—Similar to *A. muscaria*, but much smaller.
Volva—Persistent at the base of the stem, in the form of a cup; *loose and white.*	*Volva*—No cup. Base of the stem *bulbous and scaly.*	*Volva*—No cup. Base of stem not *scaly; only slightly margined by the remains of the wrapper.*
Stem—Pale yellow.	*Stem*—White.	*Stem*—White or yellow.
Gills—Pale yellow, free from stem.	*Gills*—White, rarely slightly yellow tinted.	*Gills*—Free from stem.

If the cup of a Yellow Amanita is present, the plant is safe ! If the cup is absent, it is poisonous !!

Frŏst-I-ā'-nă'



STRANGLED AMANITOPSIS (EDIBLE)

(*Amanitopsis strangulata*, Fr., Roze)

Cap greyish brown; wrapper fragments dark brown. See Genus, p. 53

GENUS CANTHARELLUS

The members of the genus *Cantharellus* differ from all other gill-bearing fungi in that the gills are in the form of shallow folds growing down the stem. The folds are generally narrow and forked or branching. The name *Cantharellus*, meaning little cup or goblet, refers to the shape of the fungus, some of the species being so deeply depressed that a cup not unlike a tall goblet results.

Cantharellus floccosus (Edible)

(See Plate Facing Page 136)

Cap or Pileus—Funnel form or trumpet shaped, depressed even into the stem. The surface of the depression is woolly or scaly. The scales sometimes persist and sometimes disappear. Reddish yellow, fading to yellow.

Gills or Lamellæ—Blunt, narrow, close; forked branched so as to appear as a network. Yellow to yellow brown.

Stem or Stipe—Short, smooth, or hairy. Sometimes extending like a root among fallen leaves.

Spores—White.

Flesh—White, thin.

Time—July to September.

Habitat—In groups in woods. The specimen pictured was found growing in moss by a stream among laurels and rhododendrons.



Section of C. floccosus

GENUS AMANITOPSIS


The genus *Amanitopsis* has white spores, the gills free from the stem, and at the base of the stem a volva, the remains of the wrapper which enclosed the young plant. *There is no annulus or ring on the stem,* a feature which separates the genus *Amanitopsis* from the genus *Amanita.*

Căn-thă-rĕl'-lŭs Flŏc-cō'-sŭs Ă-măn'-ĭ-tŏp'-sĭs

The Sheathed Amanitopsis (Edible)

Amanitopsis vaginata

Cap or Pileus—Variable in colour, ranging from white to reddish brown. Thin, fragile, smooth; or, when young, with a few fragments of the wrapper adhering to its surface. Margin distinctly and deeply striated. 2–4 inches broad.



Section of A. vaginata

Stem or Stipe—Hollow or stuffed; smooth, or sprinkled with minute mealy particles or woolly scales. Not bulbous at the base. 3–5 inches long.

Spores—White.

Ring or Annulus—None.

Wrapper or Volva—Soft, flabby, white, adhering slightly to the base.

Gills or Lamellæ—Free from the stem. White or whitish, close, irregular.

Flesh—White, or sometimes stained under the easily separating skin.

Time—June to October.

Habitat—In woods or open places, in damp vegetable mould, widely distributed. Specimen pictured was found growing in mountains of New Jersey.

A. vaginata, var. *alba*, has the whole plant white.

A. vaginata, var. *fulva*, has the cap tawny yellow.

A. vaginata, var. *livida*, has the cap leaden brown, and gills and stem tinged with smoky brown.

Văġ-ĭn-ā'-tă Fŭlv'-ă Lĭv'-ĭ-dă



SHEATHED AMANITOPSIS (EDIBLE)

(*Amanitopsis vaginata*, Roze)



HELMET MYCENA (EDIBLE)

(*Mycena galericulata*, Scop.)

Cap greyish; gills white; stems firm, hollow, hairy at the base. See p. 55

Amanitopsis parcivolvata (See Frontispiece)

Cap or Pileus—Convex, then expanded ; smooth, free from warts. Colour brilliant orange red over the whole cap; not yellow on the margin, not fading with age. Uniform in colour in young and old specimens. Margin with striations deep and long.

Gills or Lamellæ—Lemon yellow, free from stem, and rounded at the outer extremity.

Stem—Slightly tapering toward the cap; hollow. Clear lemon yellow, covered with meal-like particles.

Ring or Veil—*None* in old or young specimens.


Volva—Present; *not large and loose.* White; clinging closely to the stem in the form of scales.

Spores—White.

Flesh—Unpleasant raw odour. White, stained with orange red immediately under the skin.

Habitat—Mixed woods. New Jersey, North Carolina.

Time—July.



Section of A. parcivolvata (reduced)

The caps of the specimens found in North Carolina varied in colour from almost white to white with a red centre, and from orange or shades of orange to brilliant red.

GENUS MYCENA

The members of this genus have white spores ; no volva nor annulus ; lamellæ thin, with acute edges ; gills with a little bay cut out near the stem, sinuate, and the stem with a cartilaginous rind. Cap membranous and striate on the margin—that is, with depressed parallel lines.

Mycena hæmatopoda (See Plate Facing Page 93)

Cap or Pileus—Bell-shaped ; reddish brown, deeper shade on margin.

Gills or Lamellæ—Paler tint of colour of cap ; adnate.

Păr'-çĭ-vŏl-vā'-tă Mȳ-cē'-nă Hēm'-ă-tŏp'-ŏ-dă

Stem or Stipe—Colour of cap ; hollow ; a mere tube, with thin walls. When cut quickly, it changes to a deep red brown and exudes a purple juice. Flexible, fragile.
Spores—White.
Ring or Annulus—None.
Time—September.
Habitat—Decayed wood in forest.


GENUS LENTINUS

The genus *Lentinus* has white spores, no annulus, and no volva. The stem is central or lateral, and the lamellæ are normally toothed on their margins. The species are leathery, fleshy, and tough ; will stand drying, and revive when moist.

Lentinus lepideus is one cause of the decay of telegraph poles, railroad ties, and bridges.

Scaly Lentinus (Edible)

Lentinus lepideus



Section of L. lepideus

Cap or Pileus—Fleshy, firm, convex or expanded. Creamy white, spotted with dark brown appressed scales. 3–5 inches.
Gills or Lamellæ — Rather broad, not crowded; growing down the stem. White edges, irregularly toothed.
Stem or Stipe—Whitish. Sometimes ec-

Lĕn-tī'-nŭs Lĕp-ĭd'-ē-ŭs



SCALY LENTINUS (EDIBLE)

(*Lentinus lepideus*, Fries. Nat. size)

centric, straight, or curved; firm, solid, equal, or tapering at the base. 2–4 inches long.
Ring or Annulus—None.
Spores—White.
Flesh—White.
Time—June to August.
Habitat—On wood; common on railroad ties.

GENUS PLEUROTUS


In the genus *Pleurotus* the stem is attached to the cap at some point to one side of the centre. The stem may be on the very margin of the cap, or may be wanting altogether. The three species to be mentioned all grow on dead wood—either on dead trees or on dead branches of living trees.

The name *Pleurotus* comes from a Greek word meaning side, and has reference to the position of the stem.

Oyster Mushroom (Edible)

Pleurotus ostreatus

Cap or Pileus—Fleshy, convex, smooth, and moist. White, or tinted with ash colour or brown.
Gills or Lamellæ—White, or tinted with ash; broad; growing down the stem, and branching again and again at the base.
Stem or Stipe—Short or wanting.
Spores—*White*, oblong.
Flesh—White, tough.
Time—September.
Habitat—On dead wood. The specimens pictured were found growing on a dead forest tree at Lake Placid. The largest shells measured four inches across.




Section of P. ostreatus

Plū-rō'-tŭs Ŏs-trē-ā'-tŭs

Pleurotus sapidus (Edible)

Cap or Pileus—Convex or depressed, smooth, often irregular. White, yellowish, ashy grey, dull lilac, or even brownish; overlapping each other. 2–5 inches broad.
Gills or Lamellæ—Whitish, rather distant, growing down the stem, branching and connecting again at stem end.



Section of P. sapidus

Stem or Stipe—Stems more or less united at the base, eccentric or lateral, smooth, whitish. 1–2 inches long.
Spores—*Lilac*, oblong.
Time—June to November.
Flesh—White, tough.
Habitat—In clusters, often from a common stem, growing on decayed wood.

Elm Pleurotus (Edible)


Pleurotus ulmarius

Cap or Pileus—Convex or nearly flat, firm, smooth. White, tinted at the centre with reddish yellow or brownish yellow. 2–5 inches broad.
Gills or Lamellæ—Broad, not crowded, notched at the stem end, growing down the stem. White, turning yellow with age.
Stem or Stipe—United to the cap one side of the centre. Usually curved; solid, smooth, or downy. White or whitish. 2–4 inches long.

Săp'-ĭ-dŭs Ŭl-mā'-rĭ-ŭs



Oyster Mushroom (edible.
(Pleurotus ostreatus, Fr.)
Reduced. See page 57.



Section of P. ulmarius

Flesh—White, not tender, agreeable flavour.
Spores—White, globose.
Time—September to November.
Habitat—Conspicuous on dead parts of standing elms. Large and white. Sometimes found on other than elm trees.

GENUS HYGROPHORUS

The members of this genus may be recognized by their moist caps and by the waxy nature of their gills, which usually grow downward on the stem (decurrent), and are not very closely placed side by side. No species is known to be dangerous.

Chantarelle hygrophorus (Edible)

Hygrophorus cantharellus (See Plate Facing Page 60)

Cap or Pileus—Thin, convex at first, but later depressed. Covered with minute scales. Moist, bright red, becoming orange or yellow. ½–1 inch broad.

Hȳ-grŏph'-ŏ-rŭs

Gills or Lamellæ—Distant, somewhat arched, *growing downward on the stem* (*decurrent*). Yellow, sometimes tinged with vermilion. Unequal.
Stem or Stipe—Smooth, not truly solid, sometimes hollow. Coloured like the cap, whitish within. 2–4 inches long.
Ring or Annulus—None.
Spores—White.
Flesh—Disagreeable in flavour.
Habitat—Swamps and damp, shaded places ; in fields or woods. Common. The specimen photographed was found in dense mixed woods, Lake Placid.

Var. *H. rosea* has the cap expanded and the margin wavy.

Var. *H. flava* has the cap and stem pale yellow, the gills arched and strongly decurrent.


Var. *H. flavipes* has the cap and stem red or reddish.

Var. *H. flaviceps* has the cap yellow and the stem reddish.

Vermilion Hygrophorus (Edible)

Hygrophorus miniatus

Cap or Pileus—Thin, fragile at first, convex, becoming nearly flat. Smooth or minutely scaly. Often depressed. Red, fading to yellow or orange. 1–2 inches broad.
Gills or Lamellæ—Distant. Usually yellow, sometimes tinged with red. Notched at the stem end, or growing down the stem, or fastened to the stem by the entire width of the inner extremity of the gill.



Section of H. miniatus

Stem or Stipe—Slender, smooth. Coloured like the cap. Solid when young, hollow when old. 1–2 inches long.
Ring or Annulus—None.
Spores—White, elliptical.
Flesh—Tender, and of agreeable flavour.
Time—June to September.
Habitat—Adapts itself to varying conditions. Singly, in groups, or in clusters, in bogs or on dry hillocks. The specimen photographed was found in woods, among fallen leaves and decayed wood, Lake Placid.

Flăv'-ĭ-pēs Flăv'-ĭ-cĕps Mĭn-ĭ-ā'-tŭs



Golden-flesh Boletus. (Boletus chrysenteron, Fr.). Reduced. See page 106.



Vermilion Hygrophorus (edible). (Hygrophorus miniatus, Fr.). Reduced.



Chanterelle Hygrophorus (edible). (Hygrophorus cantharellus, Schw.)
Reduced. See page 59.



Fat Pholiota (edible). Pholiota adiposa, Fr See page 83




Honey-coloured Armillaria (edible)
(Armillaria mellea, Vahl).

Ivory Hygrophorus (Edible)

Hygrophorus eburneus (See Plate Facing Page 84)

Cap or Pileus—White, thin, fleshy, viscid.
Stem or Stipe—White, unequal, long, slender, stuffed, or hollow; viscid.
Gills or Lamellæ—White, waxy, unequal, distant, growing down the stem (decurrent), thick and firm.
Volva and Annulus—Wanting.
Spores—White.
Flesh—Odour and taste grateful.
Time—September.
Habitat—The specimen photographed was found growing among moss and fallen leaves in dense wood, Lake Placid.



Section of H. eburneus

GENUS ARMILLARIA

The members of this genus have white spores, and the gills attached by the inner extremity to the stem. The stem has a collar, but no wrapper at the base. The name is derived from the Latin *armilla*, a bracelet, referring to the ring upon the stem.


Honey-coloured Armillaria (Edible)

Armillaria mellea

Cap or Pileus—Colour from almost white to dark reddish brown. Young plants have numerous minute tufts or scales of brownish or blackish hairs. Margins sometimes striated. 1–6 inches broad.

Ĕb-ŭr′-nĕ-ŭs Är′-mĭl-lā′-rĭ-ă Mĕl′-lĕ-ă

Stem or Stipe—Usually reddish brown, paler above than below. Uniform in thickness, narrowed or slightly thickened at the base. Firm externally; soft and spongy, or hollow within. 1–6 inches long.



Section of A. mellea

Ring or Annulus—Cottony to membranous, sometimes lacking in old plants.

Gills or Lamellæ—Simply joined to the stem or running down it. White or whitish; sometimes variegated with reddish-brown spots.

Spores—White, elliptical.

Flesh—White or whitish. Taste unpleasant or acrid. Quality inferior.

Habitat—Common in woods or in cleared land, on the ground or on decayed wood. Solitary or clustered.

Time—Abundant in September. Found in June.

Var. *obscura* has cap covered with numerous small, blackish scales.

Var. *flava* has cap yellow or reddish yellow.

Var. *glabra* has cap smooth.

Var. *radicata* has tapering stem which penetrates the earth deeply.

Var. *bulbosa* has bulbous base.

Var. *exannulata* has cap smooth, margin even, stem tapering, annulus slight and evanescent, or wholly wanting.

The *Armillaria mellea* has a disagreeable taste when raw, but when cooked it is thought by some to be very good. Dr. Peck says he does not know of any unwholesome species for which it may be mistaken. The *Armillaria mellea* has the habit, very unusual for a member of the group of Agaracales, of producing from its mycelial threads tuber-like masses of fungal substances from which the fruiting caps arise. The fungal masses of the *Armillaria*, the so-called sclerotia, are ribbon or string like, and may be found between the wood and bark of cone-bearing trees. These sclerotia send out cylindrical branches, called rhizomorphs, which may penetrate the soil and attack the roots of other trees, and so continue their work of destruction in the forest.

Ŏb-scū'-rā	Glā'-brā	Būl-bō'-sā
Flāv'-ā	Răd-ĭ-cā'-tā	Ĕx-ăn'-nū-lā'-tā



GRAINY LEPIOTA

(*Lepiota granosa*, Morg.)

Cap and stem yellow-brown; gills white. See Genus, p. 63

It is to the luminosity of these mycelial threads, which permeate the decaying wood, that the weird phosphorescent light in dense woods is due.

GENUS LEPIOTA (See Plate Facing Page 64)

The members of this genus have the gills free from the stem, and have no wrapper remains at the base of the stem. In some species the cap or pileus has the surface scaly, owing to the rupture of the fibres which compose it. It is this feature which has suggested the name *Lepiota*, from the Latin word *lepis*—a scale. There are about thirty species represented in the United States, of which a few are commonly eaten.


Parasol Mushroom; Tall Lepiota (Edible)

Lepiota procera

Cap or Pileus—Convex, like an open umbrella. Thin, umbonate, covered with closely pressed scales. 3–5 inches broad.

Stem or Stipe—Long, hollow, or with cottony pith; bulbous at the base; usually covered with closely pressed scales. 5–10 inches long.

Veil or Ring—Thick and firm; often movable on the stem.



Section of L. procera

Lĕp-ĭ-ō'-tă Prōç'-ĕ-ră

Gills or Lamellæ—Closely placed, side by side. Whitish, or tinged with yellow. The inner extremity remote from the stem.
Spores—White, elliptical.
Flesh—White, soft, and dry.
Time—July to September.
Habitat—Thin woods, pastures, and by roadsides.


The specific name, *Procera*, from the Latin *procera* (tall), refers to the length of the stem.

There is no poisonous species for which it can be mistaken if one bears in mind that it has a long stem with bulbous base, a brownish, spotted cap with dark apex, and a broad basin about the insertion of the stem.

Smooth Lepiota (Edible)

Lepiota naucinoides

Cap or Pileus—Smooth, white; rarely the central part of the cap is tinged with a smoky hue. 2–4 inches broad.
Stem or Stipe—Coloured like the cap; thickened at the base. Hollow or webby. 2–3 inches long.
Veil or Annulus—White. External edge generally thicker than the inner; often movable on the stem.
Gills or Lamellæ—White when young; when old, pinkish or smoky brown. Rounded at the inner extremity and not attached to the stem. Narrower toward the stem than in the middle.
Spores—White, sub-elliptical.
Flesh—Thick, white, and tender.
Time—August—November.



Section of L. naucinoides

The smooth lepiota resembles the chalk agaric (*Agaricus cretaceous*), which has brown spores, and the meadow mush-


Nau'-çĭn-oi'-dēs



SMOOTH LEPIOTA (EDIBLE)

(*Lepiota naucinoides*, Peck)

Nat. size : Cap diam., 3½ inches; stem length, 4½ inches.



PARASOL MUSHROOM. TALL LEPIOTA (EDIBLE)

(*Lepiota procera*, Scop.)

Nat. Size : Cap Diam., 3½ inches; stem length, 4½ inches

See page 63



Lepiota Friesii, Lasch. (edible).
See page 65.



Cortinarius alboviolaceus, Fr.
Reduced. See page 86.

room (*Agaricus campestris*), which has darker gills, a persistent collar, and a stem tapering at the base. No harm can come from confusing these for edible purposes. Great care should be taken to be sure that a specimen thought to be a smooth lepiota has no volva or wrapper at the base, for the absence of a volva is the most marked difference between it and the poisonous vernal amanita (*Amanita verna*). The gills of the smooth lepiota turn a dingy brown or pink, and those of the vernal amanita remain white; and the cap of the smooth lepiota has not the moist smoothness and the brilliant whiteness of the vernal amanita.

Lepiota Friesii

Cap or Pileus—Rather thin, convex, or nearly plain, with soft, tawny fibres forming small patches over the surface. 1–4 inches wide.
Stem or Stipe—Tapering toward the cap, slightly bulbous at the base. The lower part of the stem coloured like the cap, and with similar fibrils. Hollow. Powdery white above the ring. 2–5 inches long.
Ring or Annulus—Present and pendulous.
Gills or Lamellæ—*Narrow, crowded,* free from stem; white; *some forked.*
Spores—White.
Flesh—Soft, white, with a slight odour when bruised.
Ring or Annulus—Well developed, drooping. White above, and tawny or scaly below.
Habitat—Soft, loose soil in bushy places.
Time—July to September.

GENUS MARASMIUS

The genus *Marasmius* belongs to the white-spored series. The plants are small, and wither and shrivel in dry weather, to revive again when wet. The gills are thin, and have acute edges.

The generic name comes from the same Greek word as the word marasmus, the name applied to a disease from which the patient wastes away without any apparent cause. The significance of the name will be apparent to one who watches the fleshy little plant shrink away when the sun shines.

Frē'-sĭ-ī Mă-răs'-mĭ-ŭs

Fairy-ring Mushroom (Edible)

Marasmius oreades

Cap or Pileus—Fleshy, tough, smooth, convex, or nearly plane, often with the centre higher than the space between it and the rim. Reddish ; fading, as it ages or dries, to pale yellow or buff. 1–2 inches broad.
Gills or Lamellæ—Broad and wide apart, creamy or yellowish, rounded at the stem end, unequal.
Stem or Stipe—Whitish, slender, tough, solid, coated with dense woolly hairs. 1–2½ inches long.
Ring or Annulus—None.
Spores—White.
Flesh—Thin, white, tough.
Time—May to October.
Habitat—In circles or groups.

Dangerous fungi somewhat resembling the *M. oreades*, and found in company with it, may be distinguished by their *dark-coloured spores.*

GENUS COLLYBIA

The members of this genus have white spores, and the lamellæ with thin edges attached to the stem by their inner extremity. The stem has a cartilaginous rind; that is, it is hard and of a tough texture. The genus contains fifty-four American species, some of which are regarded as edible, while others are regarded as deleterious. The velvet-stemmed collybia, or *Collybia velutipas*, is edible, and remarkable for its habit of growing long after the frosts of winter have come. It is easily recognised by its yellowish and viscid cap, and its habit of growing in tufts, and developing on the stem a *dense coat of velvety hairs.* The rooted collybia, *Collybia radicata*, may be recognised by the character of its stem, as the lower part is like a slender tap root, generally penetrating the earth to a depth equal to the length of the stem above the surface.

Ŏ-rē'-ă-dēs Cŏl-lĭb'-ĭ-ă



SPOTTED COLLYBIA (EDIBLE, Mcl.)

(*Collybia maculata*, A. & S.)

Cap and stem white with rusty spots; gills white. See Genus, p. 66



BROAD-GILLED COLLYBIA (EDIBLE, Mcl.)

(*Collybia platyphylla*, Fr.)

Surface of cap brownish, fibrillose; gills white. See Genus, p. 66




Collybia familia, Peck. Reduced (Edible)



WAXY CLITOCYBE (EDIBLE)
(*Clitocybe laccata*, Scop.)
See page 70

Collybia familia (Edible)

Cap or Pileus—Greyish, with centre darker. Smooth margin, often cracked. Slightly striate.
Gills or Lamellæ—Slightly greyish, soft, unequal, free, not crowded.
Stem or Stipe—Greyish, hollow. Lower part covered with white woolly substance.
Spores—White.
Flesh—Greenish grey.
Time—September.
Habitat—The specimen photographed was found growing upon a prostrate evergreen tree near Lake Placid.



Section of C. familia

GENUS PANUS (See Plate Facing Page 145)

The members of this genus are leathery plants, with the stems lateral or wanting. The gills are simple, not forked, and the spores are white. *Panus stypticus* is common on rotten wood, and gives out a weird, phosphorescent light.

The species of the genus *Panus* so much resemble species of the genus *Lentinus*, which do not have toothed margins, that *Panus* and *Lentinus* are considered by some as one genus, with the name *Lentinus*.

GENUS TROGIA

But one American species is reported; this is small and leathery, brownish in colour, with the spore-bearing surface white. The lamellæ are obtuse on their edges, and are not hairy. The spores are white. This plant is common on fallen branches of the alder.

GENUS SCHIZOPHYLLUM

The members of this genus have white spores and a leathery pileus, with the lamellæ hairy and grooved, or split. *Schizophyllum commune* is common on twigs or branches. It varies from ½ to 2 inches across, appearing as fluted shells on the bark.

Fă-mĭl'-ĭ-ă Pă'-nŭs Trŏg'-ĭ-ă Skĭz-ŏ-phĭl'-lŭm Cŏm-mū'-nē

GENUS OMPHALIA

The members of this genus have white spores, and the gills growing down on the stem. They have a hard, tough rind to the stem, which distinguishes them from the genus *Clitocybe*, which has fleshy stems. *Omphalia umbellifera* is a small species, with a cap about an inch broad. It is the common mushroom on the top of Mount Marcy, the highest mountain in the State of New York.


GENUS RUSSULA

The genus *Russula* may usually be recognised by its brittle character, added to its fleshy stem and the fact that the lamellæ are usually joined to the stem. Bright clear reds and purplish hues prevail, but several species exhibit a green colour, or an approach to green. The spores are white or yellowish, and the flesh never exudes a milky or coloured juice.

Emetic Russula (Dangerous)

Russula emetica

Cap or Pileus—Rosy tint to rich red. Flesh thin. The margin furrowed with parallel lines. Skin separable, somewhat viscid. Cap diameter of specimen photographed, 3½ inches.



Section of R. emetica

Gills or Lamellæ—White. Usually uniform, occasionally unequal; broad and brittle.

Stem or Stipe—Rosy. Swollen near the base. Specimen photographed, 2½ inches long.

Spores—White.

Flesh—White, unless just under the skin, where it may be pink. Very fragile. Peppery to the taste.

Ŏm-phă'-lĭ-ă Ŭm-bĕl-lĭ'-fĕ-ră Rŭs'-sŭ-lă Ĕ-mĕt'-ĭ-că



Emetic Russula (dangerous).
(Russula emetica, Schaeff.).



Green Russula (edible).
(Russula virescens, Fr.).


See page 69.

Taste—Acrid and biting.
Time—July to November.
Habitat—In damp meadows, in woods, in grassy places, under pine trees and firs.

Green Russula; Verdette (Edible)

Russula virescens

Cap or Pileus—Greyish green. At first globose, then expanded; convex or depressed at the centre. Firm and dry. Not viscid, but adorned with flaky greenish or yellowish patches, produced by the cracking of the skin. 2–4 inches broad. Margin marked with impressed lines.



Section of R. virescens

Gills or Lamellæ—White. Moderately close; free, or nearly so; narrow as they approach the stem. Some forked, others not.
Stem or Stipe—Shorter than the diameter of the cap. Smooth, white, and solid, or somewhat softer within. 1–2 inches long.
Spores—White, rough, nearly globose.
Flesh—White; mild in taste.

Vī-rĕs'-çĕns

Time—July and August.
Habitat—Grassy grounds, groves, and open woods.

No milky nor coloured juice, no coloured circular zones.

Variable Russula (Edible)

Russula heterophylla

Cap or Pileus—Variable in colour. Greenish or pinkish grey, but fleshy, firm ; slightly convex, then depressed ; smooth, and polished, the very thin skin disappearing. Margin thin, smooth, or with slightly depressed lines closely placed.
Stem or Stipe—Solid, firm, smooth, shining white, the apex occasionally dilated in the form of a cup.
Annulus or Volva—None.
Gills or Lamellæ—Narrow, *crowded*, forked ; white ; of different lengths.
Spores—White.
Flesh—White ; mild in taste.
Habitat—Woods. Common.
Time—July to October.

The specific name refers to the difference in the lengths of the gills.

GENUS CLITOCYBE

The members of the genus *Clitocybe* have the spores white, no volva nor annulus, the gills with thin edges *not* notched on the edge near the stem, and generally decurrent.

Clitocybe laccata (Edible) (See Plate Facing Page 67)

Cap or Pileus—When moist, pale red, buff red, or flesh red; when dry, greyish to pale yellowish brown. Surface covered with tiny tufts of hair. Convex when young, flattened with wavy margin when old ; often depressed. ½–2 inches broad.
Gills or Lamellæ—Broad, distant, unequal. Flesh colour. Slightly decurrent.
Veil and Annulus—None.
Stem or Stipe—Slender, stiff, fibrous, stuffed, or hollow. Colour like cap. Often twisted. 1–3 inches long.
Spores—White, rough, globose.
Flesh—Thin, pale flesh colour, leathery, tasteless.

Hĕt'ĕr-ŏ-phўl'-la. Clī-tŏ-çȳ'-bē Lăc-cā'-tă




DECEIVING CLITOCYBE (UNWHOLESOME)

(*Clitocybe illudens*, Schw.)

Dull orange; phosphorescent. See Genus, p. 70



Clitocybe virens, Scop. (edible).



Sections of C. laccata

Time—Spring to autumn.
Habitat—Wood, swamps, or open fields, naked ground, mossy or grassy places.

Var. *amethystina* has cap darker, gills amethyst, quite decurrent.

Var. *pallidifolia*, gills paler than *laccata*.

Var. *striatula*, plants small, gills showing as lines through the thin cap.

Clitocybe virens

Cap or Pileus—Fleshy, convex, expanded, obtuse. Pale greenish blue.
Gills or Lamellæ—White, crowded, thin, slightly decurrent.
Stem or Stipe—White, with occasional rusty spots ; stiff, solid. Sometimes two stems are found growing together at the base.
Spores—White.
Flesh—White.
Time—Autumn.
Habitat—The specimen photographed was found growing in mixed woods in Pennsylvania.

Ăm-ĕ-thȳs'-tĭ-nă Păl-lĭ-dĭ-fō'-lĭ-ă Strĭ-ăt'-ū-lă Vī'-rĕns


GENUS TRICHOLOMA

The members of genus *Tricholoma* have white spores, and no collar on the stem. The gills are attached to the stem, and are notched on the edge at or near the stem.

Masked Tricholoma (Edible)

Tricholoma personatum (*var. bulbosum*)

Cap or Pileus—Variable in colour; pale lilac, with yellowish tint; brighter in young specimens; sometimes whitish or pale greyish. Thick, fleshy, convex when young, with margin rolled in, and slight bloom or mealiness on the surface. When mature, smooth; margin wavy or turned upward.



Section of T. personatum (var. *bulbosum*)

Gills or Lamellæ—Faint lilac colour, with tint of violet. Narrow, unequal, free, close, and rounded at the stem end.

Stem—Faint lilac tint. Surface rather fibrous; short, stout, solid, bulbous.

Spores—Sordid white, elliptical.

Flesh—Firm.
Habitat—Thin woods, open grassy places.
Time—September to freezing weather.

Trĭ-chŏ-lō'-mă Pĕr-sŏn-ā'-tŭm



Masked Tricholoma (edible).
(Tricholoma personatum. Fr.: var. bulbosum, Pk).



Pholiota aggericola, Peck

Reduced slightly

See page 84




Craterellus cantharellus, Schw.

(Edible.) Reduced slightly

Craterellus cantharellus (Edible)

Cap or Pileus—Margin wavy. Yellow or pinkish yellow. Fleshy, firm, convex, then centrally depressed.
Gills or Lamellæ—Blunt, running down the stem, forking many times.
Stem—Solid, yellow, and smooth.
Spores—Yellowish.
Flesh—White, slightly stained in places. Taste slightly peppery; no odour.
Time—Summer and autumn.
Habitat—Specimen photographed was found growing in moss and earth on an old wood road in mixed woods, New Jersey.


Section of C. cantharellus

Craterellus is at present classified as one of the family *Thelephoraceæ*. For convenience we have placed it under "Fungi with Gills." It is interesting, as it forms a connecting link between fam. *Thelephoraceæ* and fam. *Agaricaceæ*.

GENUS AGARICUS

The genus *Agaricus* includes all brown-spored species which have free gills and a stem with a collar. The distinctive features of several edible species may be quite satisfactorily seen by reference to the table with parallel columns.

Mushroom-growing is becoming quite an important industry in this country; both professional horticulturists and amateurs successfully engage in it. It is not a difficult matter to raise the common mushroom, as the conditions necessary are easily obtained. Mushrooms will grow almost anywhere out of doors, and also in cellars, caves, and tunnels where a uniform and moderate temperature of from 50° to 60° Fahr. can be maintained. The part of a cellar devoted to mushrooms must be darkened somewhat, must have a dry floor, and must be protected from wet overhead and from winds. These conditions are common in cellars which are rather dark, but sufficiently well ventilated not to be musty. The bed for the mushrooms is prepared in a manner

Crā'-tĕr-ĕl'-lŭs Thĕl-ē-phŏ-rā'-çē-æ Ă-găr'-ĭ-cŭs

similar to that employed in making a hot-bed, care being taken that the conditions are such that too great heat is not generated. Any one desiring to experiment will find it helpful to have the "Farmers' Bulletin," No. 53, "How to Grow Mushrooms," which may be obtained from the United States Department of Agriculture, Washington, D. C.



Agaricus campestris, L. (edible).
Reduced. See page 76



Agaricus abruptus, Pk. (edible).
Reduced. See page 77.



FIELD MUSHROOM (EDIBLE)

(*Agaricus campestris*, L.)

This is the uncultivated variety of the common mushroom. See p. 76

I—PLANTS GROWING IN PASTURES, GRASSY OR OPEN PLACES

Agaricus campestris	A. Rodmani	A. arvensis	A. subrufescens
Stem—Stuffed or solid.	*Stem*—Stuffed or solid.	*Stem*—Hollow.	*Stem*—Hollow.
Gills—At first pink, later brown.	*Gills*—At first white, later brown.	*Gills*—At first slightly pink, then brown.	*Gills*—At first white, then pinkish, then brown.
Gills—About as broad as the thickness of the cap.	*Gills*—Less broad than thickness of cap.		
Collar—Simple and thin; often but traces seen in old specimens.	*Collar*—Double; a deep groove between the two layers in old specimens.	*Collar*—Double; upper part membranous, white; lower part thicker, subtomentose, radially split; yellowish.	*Collar*—Scaly on its lower surface.
Edible.	Edible.	Edible.	Edible.

II—PLANTS GROWING IN WOODS OR GROVES, OR ON SEA-COAST

A. hemorrhoidarius	A. placomyces	A. sylvaticus	A. maritimus
Flesh—Quickly changing to dull red when wounded.	*Flesh*—Not changing to dull red when wounded.	*Flesh*—Not changing to dull red when wounded.	*Flesh*—Pinkish or reddish when wounded.
Cap—White, or white with brown scales; or, when old, chocolate brown.	*Cap*—With numerous persistent brown scales.	*Cap*—Without scales, or with few evanescent ones.	*Stem*—Short or solid.
Stem—White, bulbous.			
Edible.	Edible.	Edible, but unimportant.	Edible.
Cām-pēs'-trĭs Rŏd-mān'-ī	Ăr-vĕn'-sĭs Sūb'-rū-fĕs'-çĕns	Hēm'-ŏr-rhoi-dā'-rĭ-ŭs Plā-cō'-mȳ-cēs	Sȳl-văt'-ĭ-cŭs Măr-ĭ'-tĭ-mŭs

Common Mushroom (Edible)

Agaricus campestris (See Plates Facing Pages 74 and 75)

Cap or Pileus—Silky, or with fine fibres. The young plants—"buttons"—are hemispherical or almost globose. The mature plants are convex or nearly flat. White, or tinged with dingy browns. Skin separable. Margin extending beyond the gills.

Gills or Lamellæ—Delicate pink, then dark brown, covered by a veil in young plants. Crowded, rounded at the inner extremity. Not attached to stem

Stem or Stipe—White, smooth; stuffed; that is, softer within than without. Shorter than the diameter of the cap.

Ring or Annulus—Delicate, often disappearing with age.

Flesh—White. Much esteemed.

Spores—Brownish, elliptical.


Time—Late summer and autumn.

Habitat—In pastures, lawns, and open places.

Agaricus campestris is the mushroom ordinarily seen in the markets, either fresh from the growers or preserved in cans.


Rodman's Mushroom (Edible)

Agaricus Rodmani



Sections of A. Rodmani

Căm-pĕs'-trĭs Röd-măn'-ī



RODMAN'S MUSHROOM, YOUNG (EDIBLE)

(*Agaricus Rodmani*, Peck. Reduced)



RODMAN'S MUSHROOM, MATURE (EDIBLE)

(*Agaricus Rodmani*, Peck)

Nat. size: Cap diam., 3¼ inches; stem length, 2 inches




Agaricus abruptus, Peck. (Edible)

Cap or Pileus—Creamy, with brownish spots. Firm, surface dry. Mature specimens frequently have the surface or the cap broken into large, brownish scales. 3¾ inches broad.
Gills or Lamellæ—Whitish, then pink, becoming dark brown. Narrow, close, unequal.
Stem or Stipe—Fleshy, solid, short, thick. 2 inches long.
Ring or Annulus—Double.
Spores—Brown.
Flesh—Firm, white, assuming a pinkish or salmon tint. When cut, agreeable nutty flavour.
Time—May to July.
Habitat—Grassy grounds, crevices of unused pavements. The specimen photographed was found growing in a cluster between broken stone in the gutter of a village street in New Jersey.

Agaricus abruptus (Edible)

Cap or Pileus—Creamy white, dry, and silky. Irregular in shape when young. Yellow when bruised.
Gills or Lamellæ—Slightly *pinkish* when the veil first breaks; deeper *pink* in mature specimens, becoming *brownish*. Soft, free from the stem, crowded, unequal.
Stem or Stipe—Creamy white, deeper cream, or brownish near the base. Hollow, stiff, brittle, sometimes



Section of A. abruptus

Ăb-rŭp'-tŭs

splitting lengthwise. Abruptly bulbous at the base, tapering toward cap. 3 inches long.
Spores—Brown.
Veil and Annulus—Frail, part adhering to the margin of the cap, part forming a ring on the stem.
Flesh—White, solid. Decided taste of pistachio-nut.
Time—September.
Habitat—The specimen photographed grew by the trail along Lake Placid.

GENUS HYPHOLOMA

This genus has brown spores, and no volva at the base of the stem. The veil remains as a fringe attached to the margin of the pileus, but is not always apparent in old specimens; no portion remains as a ring on the stem. The gills are attached or grown fast to the stem at their inner extremity. In this respect, and also in having the stem destitute of a collar, the genus differs from the genus *Agaricus*.

Perplexing Hypholoma (Harmless)

Hypholoma perplexum


Cap or Pileus—Convex or nearly plane, smooth, sometimes slightly umbonate. Reddish or brownish red, fading to yellow on the margins.
Stem or Stipe—Firm and hollow, slightly covered with fine threads. Whitish or yellowish to rusty red or red brown. 2–3 inches long.
Veil or Ring—Veil remains attached to the margin. No ring on the stem.
Gills or Lamellæ—Thin, crowded, slightly rounded at the stem end. At first pale yellow, then tinged with green or purplish brown.
Spores—Purplish brown, elliptical.
Flesh—Mild, not clearly bitter, white.
Time—Autumn to freezing weather.
Habitat—Grows in clusters, sometimes singly, on stumps, in the woods or in the open.

Six allied species of Hypholoma may be readily distinguished by reference to the table.

Hȳ'-phŏ-lō'-mă Pĕr-plĕx'-ŭm



Perplexing Hypholoma (harmless).
(Hypholoma perplexum, Pk.)



Section of H. perplexum

I—TASTE MILD OR NOT CLEARLY BITTER

H. epixanthus Gray-gilled Mushroom	H. capnoides	H. perplexum Perplexing Mushroom
Gills—Yellow, becoming grey; neither green nor purplish.	*Gills*—Grey to purple-brown.	*Gills*—Yellow, green, or purple brown.
Cap—Yellow or slightly tawny on disk only.	*Cap*—Yellow or slightly tawny on the disk only.	*Cap*—Brick red, yellow on the margin.
Taste—Mild.	*Taste*—Mild.	*Taste*—Mild or not clearly bitter.
		Regarded as edible.
Ĕ'-pĭ-xăn'-thŭs		Căp-noĭ'-dēs

II—TASTE BITTER


H. elæodes	H. fasciculare Tufted Yellow Mushroom	H. sublateritium Brick-red Mushroom
Gills—Green to olive.	*Gills*—Yellow to green.	*Gills*—Whitish to sooty olive.
Cap—Brick red.	*Cap*—Yellow tinged with tawny.	*Cap*—Brick red.
Stem—Hollow, rusty.	*Stem*—Hollow, yellow.	*Stem*—Solid or stuffed.
Flesh—Yellow.	*Flesh*—Yellow.	*Flesh*—Whitish.
Taste—Bitter.	*Taste*—Bitter.	*Taste*—Bitter.
	Reputed poisonous.	*Reputed poisonous and edible.*

Uncertain Hypholoma (Edible)

Hypholoma incertum

Cap or Pileus—Thin, fragile, whitish. Margin often wavy and adorned with the fragments of the woolly white veil. Opaque when dry, transparent when moist. Ovate at first, then broadly spreading. 1–2½ inches wide.

Gills or Lamellæ—Thin, narrow, close; fastened to the stem at their inner extremity. White at first, then purplish brown.



Section of H. incertum

Ĕl-æ-ō′-dēs Făsc′-ĭc-ŭ-lā′-rĕ Sŭb-lā-tĕr-ĭ′-shĭ-ŭm Ĭn-çĕr′-tŭm



Uncertain Hypholoma (edible).
(Hypholoma incertum, Pk.).



BRICK TOP (EDIBLE)

(*Hypholoma sublateritium*, Schaeff.)


Reduced. Nat. size: Cap diam., 4 inches; stem length, 4 inches

Stem or Stipe—Hollow, white, and slender. 1–3 inches long.
Spores—Purplish brown, elliptical.
Flesh—Tender.
Time—Throughout a moist season.
Habitat—In clusters, in lawns, gardens, thin woods, and pastures.

Brick Top (Edible)

Hypholoma sublateritium

Cap or Pileus—Brick red, with pale yellowish border. Surface covered with fine silky fibres. Fleshy, firm, moist. 2½–4 inches broad.
Stem or Stipe—Creamy when young, lower part slightly tinged with red. Hollow or stuffed. Silky fibres on the surface. 2½–4 inches long.
Gills or Lamellæ—Creamy when young, olive when old. Attached to stem at inner extremity. Rather narrow, crowded, unequal.
Ring or Annulus—None. Remnants of veil often seen on edge of cap.
Spores—Brown.
Flesh—Creamy, firm, bitter.
Time—September.
Habitat—The specimen photographed grew in clusters on a mossy stump in the open.



Section of H. sublateritium

GENUS STROPHARIA

There are about seven species of this genus in America. The spores are brown, there is no volva, the stem has a collar, and the lamellæ are united with the stem. They have no special economic importance.

GENUS PSATHYRA

Two American species are reported for this genus. The spores are dark brown, there is no volva, the veil is inconspicuous or wanting, and the lamellæ attached to the stem or notched at the inner extremity. The margin of the cap in the young plant and mature plant is *always straight.*

GENUS PSILOCYBE

Eleven American species are reported for this genus. The spores are dark brown. There is no volva, the veil is inconspicuous or wanting, and the lamellæ are attached to the stem or notched at the inner extremity. In the young plant the cap *curves in* at the margin.

GENUS PILOSACE

This genus has but one American species recorded. The spores are dark brown. There is no volva, and the veil is inconspicuous or wanting. The lamellæ are free from the stem.

GENUS DECONICA

But one American species is reported. This has dark brown spores, and the lamellæ growing down on the stem. The veil is inconspicuous or wanting, and there is no volva.

Strŏph-ā'-rĭ-ă Psā-thȳ'-ră Psī-lō-çȳ'-bē
Pī-lŏ-sā'-çē Dē-cŏn'-ĭ-că

GENUS CHITONIA

One American species is reported in this genus, *Clarkeinde plana*, from Nebraska. The spores are brown, and the stem has a volva at the base, but no ring.

GENUS PHOLIOTA


The members of this genus have rusty spores, and an annulus on the stem. There are about twenty known species, and some of these are edible.

Fat Pholiota (Edible)

Pholiota adiposa (See Plate Facing Page 61)

Cap or Pileus—Showy, deep yellow, with little scales of reddish brown. Fleshy, firm. At first hemispherical, then convex. Sticky when moist, shiny when dry. 2–4 inches broad.

Stem or Stipe—Stem yellow, generally rusty at the base. Equal in diameter, or slightly thickened at the base. Stuffed or solid. Tough.



Section of P. adiposa

Gills or Lamellæ—Yellowish, becoming rusty; close, and attached to the stem.

Ring or Annulus—Slightly radiating, woolly.

Spores—Rusty brown.

Time—September to November.


Habitat—In tufts, on stumps or dead trunks of trees.

Pholiota limonella, lemon-yellow pholiota, has a smaller, thinner, and more expanded cap, of a lighter yellow, with white gills.

Chĭ-tō′-nĭ-ă Phō-lĭ-ō′-tă Ăd-ĭ-pō′-să

Early Pholiota (Edible)

Pholiota præcox



Section of P. præcox

Cap or Pileus—Creamy white, smooth, not shining. Very thin skin. 1–2 inches long.
Stem or Stipe—Creamy white, rather scaly. Skin peels readily. Stuffed or hollow. 1–5 inches long.
Gills or Lamellæ—Creamy white when young, brown when mature. Soft, close, unequal, notched at the inner extremity, and attached to the stem.
Veil and Ring—Stretched like a drumhead from stem to margin of cap. Variable in manner of parting. It sometimes separates from cap margin, and forms a distinct ring about the stem; again, but little remains on the stem, and much on the rim of the cap.
Spores—Rusty brown.
Flesh—White, solid though soft, moist. Taste slightly bitter.
Time—May to July.
Habitat—Grassy ground. The specimen pictured grew about the roots of a poplar tree in New Jersey.

Pholiota aggericola

(See Plate Facing Page 73)

Cap or Pileus—Brown, shining when moist. Margin in older specimens finely and closely impressed with parallel lines. Cap diameter of specimen photographed, 2 inches.
Gills or Lamellæ—Pale brownish when young, darker brown

Prĕ'-cŏx Ăg-ġĕr-ĭc'-ō-lă



IVORY HYGROPHORUS (EDIBLE)
(*Hygrophorus eburneus*, Fries)
Reduced. Cap diam., 1 inch; stem length, 2¾ inches
See page 61



EARLY PHOLIOTA (EDIBLE)
(*Pholiota præcox*, Pers.)
Reduced. Nat. size: Cap diam., 2¼ inches; stem length, 3¼ inches.



DOG CORTINARIUS

(*Cortinarius caninus*, Fr.)

Cap lavender and brown; gills lavender; flesh lavender when bruised; taste mild; odour disagreeable

See Genus, p. 85


when mature. Attached to stem when young, free when older.

Veil—Pale brown, colour of stem.

Stem or Stipe—Pale brownish, darker at base. Softer texture within than without. Stem length of specimen photographed, 2 inches.

Spores—Rusty brown.

Flesh—Pale brownish; brittle. No taste or marked odour.



Section of P. aggericola

Time—October.

Habitat—Specimen photographed grew among grass and leaves in a New Jersey garden.


GENUS CORTINARIUS

This genus contains many species which are distinguished by the rusty yellow colour of their spores and by the webby character of the veil. It is of the utmost importance in identifying species of *Cortinarius* to know the colour of the gills of the *young plant*, as the gills of the old plants are almost uniform in colour, owing to their being dusted with the rusty yellow spores. In addition, one should carefully note the colour of the young plant compared with the old; the surface, whether sticky or dry, smooth or hairy; the taste; and the markings left on the stem by the retreating veil. All species have their gills attached to the stem, and some have them notched at the stem end. They may be looked for along the borders of woods and groves in hilly regions, during the latter part of the summer.

Cŏr-tĭ-nā'-rĭ-ŭs

Cortinarius alboviolaceus (See Plate Facing Page 65)

Cap or Pileus—Lavender, paler in the centre. Surface with rusty spots. 2½ inches broad.
Gills or Lamellæ—Crowded, unequal, attached to the stem.



C. violaceus

Stem or Stipe—Pale lavender, assuming a deeper shade when cut or bruised. Spongy in the centre, swollen toward the base. 3 inches long.
Veil—Filmy, stretching from stem to the margin of the cap in young plant.
Spores—Rusty brown.

Ăl-bō′-vī-ō-lā′-çē-ŭs




ZONED CORTINARIUS

(*Cortinarius armillatus*, A. & S., Fr.)

Cap cinnamon-brown; gills paler than cap; stem light brown with fibrous zones of red. See Genus, p. 85





THE FAWN-COLOURED PLUTEUS (EDIBLE)

(*Pluteus cervinus*, Schaeff.)

Nat. size: Cap diam., 3½ inches; stem length, 3 inches

Flesh—Tinged with violet.
Time—Autumn.
Habitat—Thick, damp woods.

C. violaceus differs from *c. alboviolaceus* in having the cap dark violet and usually covered with fibres.

GENUS PLUTEUS

The members of the genus *Pluteus* are fleshy fungi with pink spores, and gills free from the stem. They have no volva or wrapper about the young plant, and no ring or annulus on the stem. Eleven species are known from the United States, of which *Pluteus cervinus*, the fawn-coloured mushroom, is the commonest.

The generic name *Pluteus* means *all that is joined together to make a cover for besiegers at their work, that they may be screened from the missiles of the enemy.* The arrangement of the caps in the group pictured is suggestive of the meaning.


Fawn-coloured Pluteus (Edible)

Pluteus cervinus

Cap or Pileus—Light brown or fawn coloured, streaked with lines of darker brown. Surface dry and shining. Skin thin and papery. 3½ inches broad.

Gills or Lamellæ—Almost white when young, flesh colour when mature. Broad, unequal in length, free from the stem.

Stem or Stipe—Creamy white,



P. cervinus

Plū'-tĕ-ŭs Çĕr-vī'-nŭs

streaked with pale brown. Brittle, pithy in the centre when mature. 3–6 inches long.
Spores—Pink, with suggestions of yellow; salmon colour.
Flesh—White, tasteless.
Time—Early in the season, May.
Habitat—Mixed woods, on and around old stumps. New Jersey.

GENUS ENTOLOMA

The members of this genus have pink spores, and the lamellæ attached to the stem, or with a notch near the junction of gill and stem. The stem is fleshy, and not tough and hard as in *Leptonia* and *Nolanea*, genera with pink spores and adnate or sinuate lamellæ. There are some twelve species in this genus, none of which have any economic interest.

GENUS ECCILIA

The members of the genus *Eccilia* have neither volva nor annulus. The gills grow downward on the stem, the spores are pink, and the stems have a hard, tough rind, ***not fleshy as in*** *Clitopilus*. There are three species known in America.

GENUS VOLVARIA

The members of the genus *Volvaria* are fleshy fungi, soon becoming putrescent. The spores are salmon colour. A volva is present, but no annulus. Distinguished from *Amanitopsis* by having salmon-coloured spores instead of white.

GENUS CLITOPILUS

The members of this group have neither volva nor annulus. The gills grow downward on the stem, the spores are pink, and the stem is *fleshy*, without a ***hard*** and ***tough rind*** as in *Eccilia*. There are fourteen American species, of which at least two species are edible.

Ĕn-tŏ-lō′-mă	Nō-lā′-nē-ă	Ĕc-çĭl′-ĭ-ă
Lĕp-tŏ′-nĭ-ă	Vŏl-vā′-rĭ-ă	Clĭ-tŏ-pī′-lŭs



INKY COPRINUS (EDIBLE)

(*Coprinus atramentarius*, var. *silvestris*, Peck. Nat. size)

See page 91

GENUS COPRINUS

Ink Caps (Edible)

The genus *Coprinus* may be readily recognised from the fact that the spore-bearing plates dissolve to an inky fluid soon after the spores mature.

An amateur mushroom hunter may feel perfectly safe in collecting ink caps for his table, as all the species large enough to tempt the collector are not only edible, but are generally conceded to be of the best.

Their general appearance is such that even the most untrained observer should not mistake them for species of other groups.

The oblong or nearly cylindrical cap, which does not expand until ready to dissolve in inky drops, is too striking a characteristic to permit of any one making a mistake in identifying it as a specimen safe to eat.

These plants literally grow up in a night and perish in a day, as their period of growth is spent underground, and it is not until all the parts of the fruiting portions of the plants are fully developed that they push themselves above ground. Then they push and crowd from the ground in such numbers, where but a few hours before no evidence of them was seen, that each one is compressed from its cylindrical form to that of a many-sided prism, so that there would be no chance for the expansion of those within the group if it were not that those on the outer rim so rapidly expand and dissolve away.

Specimens to be eaten should be gathered in the young stage and should be cooked promptly; for though not poisonous in the black stage, they are surely not attractive.

Shelley must have had the ink caps in mind when he wrote of the fungi in the garden of "The Sensitive Plant":

> "Their mass rotted off them flake by flake,
> Till the thick stalk stuck like a murderer's stake,
> Where rags of loose flesh yet tremble on high,
> Infecting the winds that wander by."

Cō-prī'-nŭs

Shaggy-mane; Horsetail; Maned Agaric (Edible)

Coprinus comatus (See Pages i, v, viii)

Cap or Pileus—Cylindrical or barrel-shaped; becoming bell-shaped or expanded, with split margins, when old. Colour of the buttons or young plants dark; but that of the older forms white, flecked with dark patches or scales. Surface shaggy. 1½–3 inches long before expansion.

Stem or Stipe—White, smooth, hollow. 3–4 inches long.

Ring or Annulus—Slightly adherent, or movable in the young plant; later lying on the ground at the base of the stem, or wholly disappearing.

Gills or Lamellæ—Crowded. White, then tinged with pink; finally black, and dripping an inky fluid.

Spores—Black, elliptical.

Flesh—Fragile, tender, digestible, with nutty flavour.

Time—Autumn.

Habitat—Loose, rich earth. By roadsides, in pastures, and in dumping grounds.

If one study the specimens of the shaggy-mane from the time it pushes its little brown head above the ground until, as a tall black umbrella, it melts away into inky blackness, he will find much that is beautiful and interesting.

A little brown button may be cut with a sharp knife throughout its length to show the unexpanded gills lying close to the part which is afterwards to become the stem.

An older button cut in the same way will show the gills separated from the stem and the outer cover of the cap at the lower end of the gills joined to the stem. A still older specimen will show the connection of the outer cover broken loose from the base of the gill and the torn part still remaining on the stem as a temporary collar.

The outer layer of brown threads which covers the button will be found to break as the threads within expand, and to remain in the older specimens on the surface as patches of brown threads. Underlying these are broken white threads which in a younger stage, unbroken, formed a white cover under the brown. It is these loosely hanging threads which give the shaggy appearance to the cap of the mature plants and which have suggested the names of shaggy-mane, horsetails, and comatus (*comatus*, in Latin, meaning hairy).

Cō-mā'-tŭs



INKY COPRINUS (EDIBLE)

(*Coprinus atramentarius*, Fries. Nat. size)

Inky Coprinus (Edible)

Coprinus atramentarius

Cap or Pileus—Cylindrical at first, broadening by degrees until it is cone-shaped. Colour greyish or greyish brown, with suggestions of lead colour. Smooth or with a few obscure scales on the disk. Often suffused with bloom. The margin sometimes notched or lobed. Deliquescing. 1–3 inches in diameter.
Stem or Stipe—Slender, smooth, whitish, hollow.
Ring or Annulus—A slight vestige of one may be seen to extend around the stem near the base as an irregular zigzag elevated line of threads.
Gills or Lamellæ—Crowded. At first whitish and flocculose on the edges, then black, moist, dropping away in inky fluid.
Spores—Black, elliptical.
Flesh—White, quickly deliquescing.
Time—Autumn.
Habitat—Rich soil, waste places, woods.

The form growing in the woods is much more beautiful and is known as *C. atramentarius*, var. *silvestris*. See plate facing page 89.

Glistening Coprinus (Edible)

Coprinus micaceus

Cap or Pileus—Ovate, then bell-shaped. Striations radiating from near the centre of the disk to the margin. Glistening mica-like scales cover undisturbed young specimens. 1–2 inches broad. Colour tan, light buff, or tawny yellow.
Stem or Stipe—Slender, smooth, fragile, white, hollow. 1–3 inches long.
Ring or Annulus—Rarely seen except in very young specimens.
Gills or Lamellæ—Not as crowded as in the ink cap and shaggy-mane. Colour white, then tinged with pinkish or purplish brown, finally black.
Spores—Brown, elliptical.
Flesh—A nutty flavour when raw. In wet weather it melts to an inky fluid. In dry weather it may dry with all parts well preserved.
Time—Common during spring and early autumn.

The glistening coprinus is small and beautiful, and grows in clusters on decaying woods, stumps, or buried roots.

Ăt'-ră-mĕn-tā'-rĭ-ŭs Mĭ-cā'-çē-ŭs

GENUS GOMPHIDIUS

This genus has black spores. The lamellæ are waxy, and grow downward on the stem.

GENUS PSATHYRELLA

The members of this genus are fleshy fungi, with black spores. They have no annulus, and the lamellæ are not waxy or decurrent. The cap is striate, and the stem has a hard, tough rind.

GENUS PANÆOLUS

There are six or more species of this genus in America. They have black ovoid spores, a cap smooth and not striate, a fleshy stem.

GENUS LACTARIUS

A milky or coloured juice exuding from the broken gills of a fungus marks it as a *Lactarius*. The species are mostly stout and fleshy, the cap resembling an inverted cone; the gills are more or less decurrent, and the stem short and stout. Many of the species have a hot, acrid taste, and some have zones of colour on the surface of the cap. The spores are white or yellow. The juice exuded may be white or orange or blue.

Peppery Lactarius (Edible)

Lactarius piperatus

Cap or Pileus—Creamy white, fleshy, firm, depressed in the centre. Dry, never viscid, and uncommonly broad. 3–12 inches in diameter.

Gills or Lamellæ—Creamy white, narrow, crowded, unequal, forked, decurrent. Exuding a milky juice when bruised. *Milk white.*

Stem or Stipe—Creamy white, short, thick, solid, rounded at and slightly tapering toward the base.

Gŏm-phĭd'-ĭ-ŭs Sā-thȳ-rĕl'-lă Păn-æ'-ō-lŭs
Lăc-tā'-rĭ-ŭs Pĭp-ĕr-ā'-tŭs



PEPPERY LACTARIUS

(*Lactarius piperatus*, Scop. Nat. size)



PEPPERY LACTARIUS

(*Lactarius piperatus*, Scop. Nat. size)




Mycena hæmatopoda, Pers.

See page 55



Lactarius gniotus, Fr.



L. piperatus

Veil and Annulus—None.
Spores—White.
Flesh—Slightly paler than the surface of the cap. Taste very peppery. Odour quite aromatic.
Time—Summer.
Habitat—Specimen pictured was found in mixed woods, among dried leaves and sticks, in New Jersey.

Lactarius ligniotus

Cap or Pileus—Fleshy; brown velvety surface; convex to plane; brittle.
Gills or Lamellæ—Attached to the stem, distant, unequal. Snow white to yellowish white. *Milk mild and white.*
Stem and Stipe—The upper part the same colour as the cap surface, base lighter; hollow.
Ring or Annulus—None.
Spores—White.
Flesh—Taste pleasant. Broken flesh reddish white, then yellowish.
Time—September.
Habitat—Common among mosses under fir trees. The specimen pictured was found growing in mixed woods at Lake Placid.

Lĭg-nĭ-ō'-tŭs

CHAPTER VI: FUNGI WITH TEETH—HYDNACEÆ

THE fungi with teeth are so called because, instead of bearing their spores on the surface of gills and pores, they bear them on the surface of awl-shaped teeth, which project downward. The genera of the family *Hydnaceæ* are distinguished by the size, shape, and attachment of the teeth. Plants with teeth only, and no basal membrane, make the genus *Mucronella.* Plants with flattened, leaf-like teeth attached to a leathery membrane, growing on wood, either in the form of a cap, or simply spreading over the host, make the genus *Irpex.* Plants with thick, blunt, irregular spines make the genus *Radulum.* Fleshy or membranous plants with caps and flattened teeth, growing on the ground, make the genus *Sistotrema.* Plants which spread over their host, closely attached to its surface, and have simple, bristle-like teeth, make the genus *Pycnodon.* Plants growing in a manner similar to those of the genus *Pycnodon,* but having low-crested wrinkles instead of bristles, make the genus *Phlebia;* while those with smooth hemispherical warts make the genus *Grandinia,* and those with crested papillose warts make the genus *Odontium.* The typical genus *Hydnum* has the teeth cylindrical, so that a cross section would appear circular. This is the only large genus, and in it are found several important edible species. These may be put in two groups, one containing the species with a cap and central stem, and one the species growing in branched masses with no distinct cap. These are commonly known as Hedgehog Mushrooms.

Hȳd-nā'-çĕ-æ Ȳr'-pĕx Sȳs-tō-trē'-mă Phlĕb-ȳ'-ă
Mū-crō-nĕl'-lă Răd'-ū-lŭm Pȳc'-nō-dŏn Grăn-dȳ'-nȳ-a
Ō-dŏnt'-ȳ-ŭm Hȳd'-nŭm

GENUS HYDNUM

Spreading Hydnum (Edible)

Hydnum repandum (See Plate Facing Page 103)

Cap or Pileus—Fleshy, fragile, moist, smooth or somewhat scaly in mature specimens. Variable in colour; light red, pale buff, or rusty yellow. Convex, plane, or deeply concave by the stem becoming hollow. Margins often wavy. 1–4 inches broad.

Teeth or Aculei—Pointed, whitish, easily detached, leaving little cavities in the fleshy cap.


Stem or Stipe—Stem solid in young specimens, hollow in older specimens. Surface rather rough; often eccentric. 1–3 inches long.

Flesh—Watery, lighter colour than cap.

Spores—Whitish.

Time—July to October.

Habitat—Woods and open places. Singly or in clusters. The specimen pictured was found in North Carolina in February.



Section of H. repandum

Hydnum rufescens (edible) is redder than the typical form, is smaller and more regular.

White Hydnum (Edible)

Hydnum albidum

Cap or Pileus—White, fleshy. 1–2 inches broad.
Teeth or Aculei—White, short.
Stem or Stipe—White, solid, short. 1–2 inches long.
Flesh—White.
Time—June to August.
Habitat—Thin, wet woods.

Rē-păn'-dŭm Rū-fĕs'-çĕns Ăl-bī'-dŭm

Hydnum imbricatum



H. imbricatum

Cap or Pileus—Brownish, as if scorched. Surface cracked in irregular scales. 6–7 inches wide.
Teeth—Bluish grey.
Stem or Stipe—Short and thick, with irregular scales.
Flesh—Whitish.

Time—Late summer.
Habitat—Dry woods.

Bear's-head Hydnum (Edible)

Hydnum caput-ursi

Plant—White, fleshy.
Branches and Teeth—Short branches covered with awl-shaped teeth of varying length, pointed toward the ground.




Section of H. caput-ursi

Flesh—White when young, creamy when old.
Habitat—On prostrate or standing tree trunks of decaying deciduous trees. Erect, if on the upper side; ascending or pendulous or both, if on the side of the trunk.

The bear's-head hydnum is very variable in form. The

Ĭm-brĭ-cā'-tŭm Că'-pŭt-ŭr'-sī



BEAR'S HEAD (EDIBLE)

(Hydnum Caput-Ursi, **Fries)**



CORAL HYDNUM

(*Hydnum coralloides*, Scop.). See p. 97

essential character is a solid, fleshy body with short protecting branches bearing numerous simple or branched teeth, which may vary in length from one-sixth of an inch to two inches.

Medusa's Head (Edible)

Hydnum caput-medusæ

Hydnum caput-medusæ has knob-like branches, with short, distorted teeth above, and long, uniform ones below. At first it is white, but later changes to a smoky or ash colour. The change in colour from white to an ashy tint distinguishes it from *H. caput-ursi*, which changes to creamy white.

Hedgehog Hydnum (Edible)

Hydnum echinaceus

The hedgehog hydnum is white at first, and then creamy. The branches are knob-like, and bear numerous straight, equal teeth about two inches long.

Coral Hydnum (Edible)

Hydnum coralloides

Plant—Pure white, becoming creamy with age.
Branches and Teeth—Numerous, spreading, angular or flattened. Terminal branches often curved upwards, terminating in a crowded mass of spines. Teeth, 1/6–1/3 of an inch long.
Stem—Short, dividing into branches almost from the base.
Flesh—Tender, white, with agreeable taste.
Time—July to October.
Habitat—On prostrate trees in mountainous or hilly country.

When a mere child, Elias Fries was so attracted by this beautiful coral-like fungus, which grew in his native forests in Sweden, that he was led to take up the study of fungi, and later became one of the most prominent students in that branch of botany, and laid the foundation for the study of the *Basidiomycetes*.

Că'-pŭt-mē-dū'-sæ Ĕ͡ch-ī-nā'-çē-ŭs Cŏr-ăl-loi'-dēs

CHAPTER VII: FAIRY CLUBS AND CORAL FUNGI—CLAVARIACEÆ

THE fairy clubs and coral fungi belong to the family *Clavariaceæ*. They are fleshy fungi of upright growth, which have their spore-bearing surface exposed on the apices of branching or simple club-like forms. Many are extremely beautiful, resembling corals of exquisite shades of pink, violet, yellow, or white.

The seven genera are distinguished by the colour of the spores, by their habit of growth—whether simple or branched, and, if branching, by the form of the branches; whether club-like or thread-like, flat or round, cartilaginous or leathery. Many of the members of this family are edible, and none are known to be unwholesome, so that it will be safe for a beginner to try any of them.

GENUS PHYSALACRIA

Plant small, simple, hollow, and enlarged at the apex.

GENUS PISTILLARIA

Plants conspicuous, club-shaped or thread-like, with two spicules to each spore-bearing cell.

GENUS TYPHULA

Plants conspicuous, club-shaped or thread-like, with four spicules to each spore-bearing cell.

Clăv'-ār'-ĭ-ā'-çē-æ Phўs-ă-lā'-crĭ-ă Pĭs-tĭl-lā'-rĭ-ă Tўph'-ū-lă



LITTLE TONGUE CLAVARIA

(*Clavaria ligula*, Fr.)

Young specimens yellow; mature specimens tawny. See p. 99

GENUS SPARASSIS

Plants conspicuous, with the branches strongly flattened or leaf-like.

Sparassis crispa somewhat resembles a yellow cauliflower, and often forms masses as large as one's hand. It is considered an excellent fungus for the table.

GENUS PTERULA

The members of this genus are mostly slender, thread-like forms, cartilaginous when moist, and horny when dry.

GENUS LACHNOCLADIUM

The members of this genus are leathery plants covered with hairs.

GENUS CLAVARIA

The genus *Clavaria* is the largest genus in the family *Clavariaceæ*. Many of the species are edible, and so easily recognised that the beginner may venture without hesitation to eat any of the branching forms. The club-like forms might be mistaken for certain club-shaped forms of the sac-fungi unless an examination of the spores were made. The *Clavarias* would have the spores on little spicules, as in the garden mushroom, whereas the forms for which they might be mistaken would have them in membranous sacs.

In collecting species of *Clavaria*, notes should be taken as to the character of the apices of the branches, the colour of the branches, the colour of the spores, the taste, and the place of growth.

Pale Yellow Clavaria (Edible)

Clavaria flava

Plant—White and pale yellow. 2–5 inches high; the mass of branches 2–5 inches wide.

Spā-rās'-sĭs Crĭs'-pă Tĕr'-ŭ-lă Lăk-nŏ-clā'-dĭ-ŭm Clăv-ăr'-ĭ-ă Flāv'-ă

Branches—Round, not flattened; smooth, crowded, and nearly parallel, pointing upward. Whitish or yellowish, with pale yellow tips. Branchlets terminating in from one to three blunt, tooth-like points.
Stem—Short, thick, white.
Spores—Yellowish.
Flesh—White, tender.
Time—July to September.
Habitat—Thin woods and open places.

Golden Clavaria (Edible)

Clavaria aurea

The plant is from three to four inches high, with the branches of a uniform deep golden yellow, and often longitudinally wrinkled. The stem is stout, but thinner than the *Clavaria flava*, which it somewhat resembles.

Red-tipped Clavaria (Edible)

Clavaria botrytes

Plant—From 2–5 inches high, whitish or yellow or pinkish, with the *tips of the branches red.*
Branches — Sometimes longitudinally wrinkled, repeatedly branched.
Stem—Short, thick, fleshy, whitish.
Time—July to September.
Habitat—Thin woods and open places.

Crested Clavaria (Edible)

Clavaria cristata

Plant—Small, not more than 2–2½ inches high. White or whitish, often faintly tinged with dull pink, or creamy yellow, or smoky tints.
Branches—Widened and flattened above, and deeply cut into several finger-like points, which may turn blackish brown when old.
Stem—Slender, spongy within.
Spores—White.
Habitat—Woods and open places, especially in cool, shady, moist places.

Au'-rĕ-ă Bō-trȳ'-tēs Crĭs-tā'-tă



Golden Clavaria (edible).
(Clavaria aurea, Schaeff.).



Clavaria formosa, Pers. (edible).

See page 101.

Pistil Clavaria; Large Club (Edible)

Clavaria pistillaris

Of the club-shaped clavarias, the pistil clavaria is the largest. It is of a light yellow tinged with brown or red, and with soft white flesh. In shape it resembles an Indian club, being blunt and rounded at the summit, with a diameter of an inch or less, and a height of five inches or less. It is found during the summer in grassy open places or in thin woods.

Clavaria fellea

The clubs of this species are about one inch high, light yellow tinged with brown or red, somewhat divided by pairs from bottom to top into two forked branches. The stem is round and solid, and the branches are crowded and nearly parallel, with blunt tips, and of a uniform colour. The taste is bitter.

Clavaria formosa

The *Clavaria formosa* has a stout whitish stem, with erect branches, dividing and subdividing repeatedly, golden to pink, the branchlets obtuse. The specimen pictured grew on a fallen tree in dense mixed woods.

Pĭs'-tĭl-lā'-rĭs Fĕl'-lĕ-ă Fŏr-mō'-să

CHAPTER VIII: FUNGI WITH PORES—BOLETACEÆ; POLYPORACEÆ

THE fungi with pores naturally divide into two groups. The perishable fleshy fungi with *pores easily separating* from the cap and from each other make the family *Boletaceæ*. The perishable fleshy fungi, and the leathery, corky, membranous or woody fungi, with *pores permanently united* to the cap and to each other, make up the family *Polyporaceæ*. In each the spores are borne on the linings of pores or tubes placed close together, so that on the surface of the cap they look much like large pin pricks on a stiff cushion; but in a sectional view, obtained by cutting the cap from the upper surface to the lower, they look like tubes placed side by side.



Section of a Boletus

Among the fleshy forms the genus *Fistulina* has the stem lateral and the tubes close together, but distinct from each other. The genus *Boletus* has the tubes easily separable from the cap, and the genera *Boletinus* and *Strobilomyces* have the tubes quite firmly connected. A brief comparison of essential points is given in the table.

1. Stem strictly lateral. Mouths of the tubes separated from each other..... Genus FISTULINA
2. Stem central. Pores readily separating from the cap..... Genus BOLETUS
3. Stem central or eccentric; fleshy. Pores in more or less radiating rows..... Genus BOLETINUS
4. Stem central, *tough*. Pores uniform. Cap woody..... Genus STROBILOMYCES

GENUS FISTULINA

The genus *Fistulina* contains one notable species, *Fistulina hepatica*, so called from its resemblance to a liver. In its early stages it somewhat resembles a strawberry, and later it may

Bō-lĕ-tā'-çĕ-æ	Fĭs-tū-lī'-nă	Bŏ-lĕ-tī'-nŭs
Pō'-lў-pŏ-rā'-çĕ-æ	Bŏ-lē'-tŭs	Strō-bĭl-ō'-mў-çēs



CONE-LIKE BOLETUS (EDIBLE, Mcl.)

(*Strobilomyces strobilaceus*, Berk.)

Purplish black; flesh white, changing to red when bruised. See p. 102



Painted Boletinus (edible).
(Boletinus pictus, Pk.).



Spreading Hydnum (edible).
(Hydnum repandum, L.).
See page 95.

have the appearance of a big red tongue protruding from a tree trunk, so that the French call it *langue de bœuf.* It is often called vegetable beefsteak, from the flesh-like fibre and colour of the fresh specimens. When young, the upper side is velvety and of a fine peach colour ; later it becomes liver red and loses the velvety appearance ; the under surface is flesh coloured, and is rough, much like the surface of a tongue, owing to the fact that the tubes are free from one another.

GENUS BOLETINUS

The spore-bearing surface of the genus *Boletinus* is composed of broad, radiating lamellæ connected by numerous narrow partitions so as to form large angular pores. The tubes are not easily separable from each other and from the portion on which they are borne. They are yellowish in colour, and grow downward somewhat on the stem. The genera may be distinguished by the following table:

1. Stem hollow..... BOLETINUS CAVIPES
2. Stem solid, lateral, or eccentric..... BOLETINUS POROSUS
3. Stem solid, central. Cap pale yellow and silky..... BOLETINUS DECIPIENS
4. Stem solid, central. Cap red..... BOLETINUS PALUSTER
5. Stem solid, central. Cap with red scales..... BOLETINUS PICTUS

Painted Boletinus (Edible)

Boletinus pictus

Cap or Pileus—Convex or nearly plane; at first covered with *red matted fibres, which soon divide into small scales,* so as to reveal the yellow colour of the cap beneath. 2–4 inches wide.
Tubes—Pale yellow, or pale yellow tinged with brown. Not easily separated from the cap.
Veil—Webby, concealing the tubes of the young plant.
Stem or Stipe—Solid, with scales and colour similar to those of the cap. 1½–3 inches long.
Ring or Annulus.—Webby; evident in some specimens.
Spores—Pale yellow tinged with brown.
Flesh—Yellowish, often assuming reddish tints when bruised.
Habitat—Woods and mossy swamps.

Căv'-ĭ-pēs Pō-rō'-sŭs Dē-çĭp'-ĭ-ĕns Pă-lŭs'-tĕr Pĭc'-tŭs

GENUS BOLETUS

The species in the genus *Boletus* are numerous, and many are extremely beautiful. They are distinguished from the other pore-bearing fungi by the fact that their tubes are easily separable from each other and from the portion on which they are borne.

They are to be looked for in the warmest part of the season, and with a few exceptions will be found on the ground. As it is difficult to dry specimens so that they will retain their size, shape, and colour, careful notes should be taken of these points, together with the colour of the spores, the colour of the tubes, the colour of the flesh before and after being bruised, the character of the stem, the presence or absence of hairs and fibres on the several parts of the plant. The genus *Boletus* contains many edible species, and also many which are dangerous; and as the recognition of the different species is a difficult matter, even for experts, the beginner should be especially cautious. In general it will be well to avoid for edible purposes all *Boleti* which *change colour* on exposure to the air or on being pressed; all those which have *red-mouthed tubes*, or mouths of a deeper colour than the rest of the tube; and all those with a *peppery* or *acrid* taste. Experiment cautiously with the other species.

A detailed description of the species is out of the question in this book; for such the reader must refer to "Boleti of the United States,"* or to "Führer für Pilzfreunde."†

Boletus glabellus

Cap—Smoky yellow.
Flesh—White, *changing to blue when wounded.*
Tubes—Brownish yellow tinged with green, *changing to blue when wounded.*
Stem—Reddish at the base, pallid above, with a narrow reddish *circumscribing zone or line at the top.*
Habitat—Grassy ground.

Glā-bĕl'-lŭs

* Bulletin, New York State Museum, No. 8, 1888. This can be obtained at a small price of the State Librarian at Albany, New York. In it are described one hundred and ten species.

† By Edmund Michael. A book in German, with sixty-eight coloured plates.

Boletus bicolor

Cap—Dark red, fading when old ; often marked with yellow.
Flesh—Yellow, slowly *changing to blue* when wounded.
Tubes—Bright yellow, *changing to blue.*
Stem—Solid, *red, generally yellow at the top.*
Habitat—Woods and open places.

Boletus cyanescens

Cap—Pale buff or greyish yellow. Covered with woolly scales, two-fifths of an inch wide.
Flesh—White, *quickly changing to blue* when wounded.
Tubes—White, becoming yellowish, *changing to blue* when bruised.
Stem—Coloured like the cap. Swollen, stuffed, not covered with a network.
Habitat—Woods and open places.

Boletus pallidus

Cap—Pallid or brownish white, sometimes tinged with red.
Flesh—White.
Tubes—Pale whitish yellow, *changing to blue* when wounded.
Stem—Long and whitish, sometimes streaked with brown ; smooth.

Boletus mutabilis

Cap—*Brown*, becoming blackish ; smooth and shining ; or dry, somewhat woolly.
Flesh—Bright yellow, *promptly changing to blue* when wounded.
Tubes—Yellow, changing to greenish yellow, and *quickly becoming blue when wounded.* Mouths large and angular.
Stem—Stout, bright yellow within, covered with *brown and dotted scales.*
Habitat—Dense woods.

Boletus speciosus

Cap—Red. 3–7 inches broad.
Flesh—Pale or bright lemon yellow, *changing to blue* when wounded.

Bĭ'-cō-lŏr Çȳ-ăn-ĕs'-çĕns Păl'-lĭ-dŭs Mū-tă'-bĭ-lĭs Spē-çĭ-ō'-sŭs

Tubes—Bright lemon yellow, becoming dingy yellow with age; *changing to green, then to blue,* when wounded.
Stem—*Bright lemon yellow without and within,* sometimes reddish at the base. 2–4 inches long.
Habitat—Thin woods.

Golden-flesh Boletus

Boletus chrysenteron (See Plate Facing Page 60)

Cap—Yellowish brown, reddish brown, brick red, or olive tinted with reddish chinks.
Flesh—Yellow, *red just under the skin,* often changing to blue when wounded.
Tubes—Greenish yellow, *changing to blue when wounded.*
Stem—Red or pale yellow.
Habitat—Woods or mossy banks, common.

Boletus radicans

Cap—Dry, somewhat woolly. Greenish grey, becoming pale yellow. Margin rolled under.
Flesh—Pale yellow, instantly *changing to dark blue* when wounded.
Tubes—Lemon yellow.
Stem—Tapering downward and rooting. Woolly, with a reddish bloom. Pale yellow, becoming dark with a touch.
Habitat—Woods, Ohio.

Boletus Peckii

Cap—Firm, dry. *Red, fading to yellowish red or buff brown.*
Tubes—Yellow, *changing to blue* when wounded.
Stem—Red; yellow at the top.
Habitat—Woods.

Boletus calopus

Cap—Olive tinted, somewhat woolly. 2–3 inches wide.
Flesh—Pallid, *slightly changing to blue* when wounded.
Tubes—Yellow.
Stem—Covered with a network. Wholly scarlet, or at the apex only. Longer than the diameter of the cap.
Habitat—Woods.

Crĭs-ĕn'-tĕr-ŏn Rā'-dĭ-cănns Pĕck'-I-I Căl-ō'-pŭs



BITTER BOLETUS (UNPALATABLE)

(*Boletus felleus*, Bull.; var. *obesus*, Pk.). See p. 107


Purple Boletus

Boletus purpureus

Cap—Dry, *velvety. Purple red.*
Flesh—*Changing to blue,* and then to dark yellow in the young plant.
Tubes—Yellow or greenish yellow. Mouths minute, *orange tinged with purple, changing to blue* where wounded.
Stem—Stout, with purple veins or dots. Apex sometimes covered with a network. Yellow, reddish within.
Habitat—Woods.

Boletus Satanus

Cap—*Smooth,* somewhat sticky. *Pale brown or whitish.* 3–8 inches broad.
Flesh—Whitish, *becoming reddish or violet* when wounded. *Very poisonous.*
Tubes—Yellow. Mouths bright-red, becoming orange.
Stem—Thick, swollen, marked above with a red network. 2–3 inches long.
Habitat—Woods.



B. Satanus

Bitter Boletus

Boletus felleus

Cap—Smooth, pale yellowish, greyish brown, yellowish brown, reddish brown, or chestnut. 3–8 inches broad.
Flesh—White, changing to flesh colour when bruised. Taste *bitter.*
Tubes—White, becoming flesh colour. Mouths angular. Adnate to the stem.
Stem—Covered with network. Colour similar to the cap. 2–4 inches long.
Habitat—Woods and open places.

Pŭr-pū'-rĕ-ŭs Să-tā'-nŭs Fĕl'-lĕ-ŭs

Boletus scaber

Cap—*Smooth, viscid when moist,* or minutely woolly, velvety, or scaly. 1–5 inches wide. Colour ranges from nearly white to almost black.
Tubes—Free from the stem, white, long. Mouths minute.
Flesh—White.
Stem—Long, ground colour whitish, *roughened with blackish brown or reddish dots or scales.* 3–5 inches long.
Habitat—Common in woods, swampy and open places.

Orange-cap Boletus

Boletus versipellis

Cap—Orange red. 2–6 inches wide. Dry, minutely woolly, then scaly or smooth. Margin edged with the remains of the veil.
Flesh—White or greyish.
Tubes—Long. Mouths minute. Greyish-white.
Stem—Solid; white scaly wrinkles. Whitish colour. 3–5 inches long.
Habitat—Woods and open places.

Chestnut Boletus

Boletus castaneus

Cap—Dry, *minutely velvety. Cinnamon or reddish-brown.* 1–3 inches broad.
Flesh—White.
Tubes—White, becoming yellow. Short, and free from the stem.
Stem—Clothed and coloured like the cap. 1–2½ inches high.
Habitat—Woods and open places.

Boletus eximius

Cap—*Purplish-brown or chocolate colour,* fading to smoky red or pale chestnut. 3–10 inches broad.
Flesh—Greyish or reddish white.
Tubes—Resembling the cap in colour. Mouths minute.
Stem—Stout, *covered with a meal-like powder.* Colour resembling the cap, grey tinged with purple within. 2–4 inches long.
Habitat—Woods.

Scā'-bĕr Vĕr-sĭ-pĕl'-lĭs Căs-tā'-nĕ-ŭs Ĕx-ĭm'-ĭ-ŭs



SCABROUS-STEMMED BOLETUS (EDIBLE)

(*Boletus scaber*, Fr.; var. *niveus*, Gill.)

Cap white. See p. 108

Edible Boletus

Boletus edulis

Cap—Convex or nearly plane; *smooth*, moist; compact, then soft. Greyish red, brownish red, or tawny brown. 4–6 inches broad.
Flesh—White or yellowish; reddish beneath the skin.
Tubes—Convex, nearly free, long, minute, round. White, then yellow and greenish.
Stem—Short or long, straight or curving, sometimes bulbous, stout, covered with network. Just beneath the stem whitish or brownish. 2–6 inches long.
Habitat—Woods and open places.

Boletus subtomentosus

Cap—*Covered with soft woolly hairs.* Somewhat olive green, uniform in colour under the skin, yellow chinks on the surface.
Flesh—White.
Tubes—Yellow, with large angular mouths.
Stem—Stout, rugged, with minute dots.
Habitat—Common in woods.

Boletus Americanus

Cap—Thin, soft, viscid, *slightly woolly on the margin when young.* Yellow, becoming dingy with age; sometimes streaked with bright red. 1–3 inches broad.
Flesh—Pale yellow.
Tubes—Not free from the stem. Large, angular. Pale yellow, becoming tinged with brown.
Stem—Slender. No annulus. Yellow, brownish towards the base, marked with numerous brown or reddish-brown glandular dots; yellow within. 1½–2½ inches long.
Habitat—Woods, swamps.

POLYPORACEÆ

The fungi with pores permanently united to the surrounding tissue and to each other form a large and important group, the *Polyporaceæ*. With but few exceptions they are leathery, corky, membranous, or woody.

Nearly six hundred species have been reported from America.

Ĕd'-ū-lĭs Sŭb'-tō-mĕn-tō'-sŭs Ă-mĕr'-ĭ-cā'-nŭs Pŏ'-lĭ-pŏ-rā'-çē-æ

GENUS MERULIUS

Merulius lacrymans

The simplest of these *Polyporaceæ* is the dry-rot fungus, *Merulius lacrymans*. The food-seeking portion consists of fine white threads, mycelium, which penetrate the woodwork of buildings, causing it to crumble to dust. The fruiting portion consists of flat, irregular bodies whose under surface bears the spores in shallow pits separated by narrow ridges. Water is excreted from these flat disks, which, from the habit of dropping like tears, has suggested the specific name *lacrymans*, from the Latin *lacrymare*, to weep.

GENUS POLYPORUS

The genus *Polyporus* and the genus *Trametes* have the pores closely packed and united to together. In *Trametes* the uniting substance is the same as the substance of the cap, but in *Polyporus* the uniting substance is different from the substance of the cap. The genus *Polyporus* in its widest sense is a large, one, numbering some five hundred species, and containing every texture from fleshy or pulpy to woody.

The fruiting portion is often seen in the form of brackets shelving out from standing or fallen trees. The mycelium penetrates the wood, softening it, and causing it to crumble, so that in the course of time the tree dies. The external evidence that the tree has been attacked is the appearance of the fruiting portion, which often attains a very great size.

Elfvingia (polyporus) megaloma

Polyporus megaloma is attractive to frequenters of the woods, as its spore surface when fresh is soft and white, and furnishes an attractive surface for stencil drawings.

Elfvingia (polyporus) fomentaria

The brackets of *Elfvingia fomentaria* resemble horses' hoofs. They are employed in the manufacture of *amadou*, or

Měr-ū'-lĭ-ŭs Lăc'-rў-măns Pŏ-lўp'-ŏ-rŭs Trā-mē'-tēs Měg-ă-lō'-mă Fō-měn-tā'-rĭ-ŭs



TINDERWOOD POLYPORUS

(*Elfingia fomentaria*, L.; *Fomes fomentarius*, Gill.; *Polyporus fomentarius*, Fr.)
Upper surface grey to brown; tube-surface concave. See p. 110

German tinder, which, in the form of sticks or fusees, commonly known as punk, is used for lighting cigars and pipes in the wind, or for touching off fireworks.

The fusees are made by beating the fungus substance until it is flexible, and then dipping it into saltpetre,

Polyporus conchatus

Polyporus conchatus is a beautiful species found on the under side of fallen trees, growing in such a manner that from the spore-bearing surface the shells look like rosettes of shimmering golden-brown velvet.

Polyporus velutinus

Polyporus velutinus is common in the form of stiff ruffles with white or creamy spore surface, and grey or tan-coloured plush-like upper surface.

Polyporus pergamenus

Polyporus pergamenus has a somewhat bristle-like, tan-coloured spore surface, and a velvety drab upper surface, often tinged with green from a green alga which grows upon it.

Polyporus perennis

Polyporus perennis resembles a slender-stemmed goblet.

Polyporus sulphureus

Polyporus sulphureus is an edible fleshy form. It grows in clusters, the stemless caps often five or six inches broad and overlapping. The colour of the young cap is yellowish red or pale orange, with the margin wavy, and of a beautiful yellow. It is soft and juicy, and often—when cut—exudes a yellow juice.

The plants may be found on the dead wood of trees during showery weather from May to October. The mature plants become dry and crumbling, and as they dry they lose

Cŏn-chă'-tŭs Vĕl-ū'-tĭ-nŭs Pĕr-gă-mĕ'-nŭs Pĕr-ĕn'-nĭs Sŭl-phū'-rĕ-ŭs

their attractive colours. They exhibit phosphorescence in early stages of decay. The tubes are minute and short and of a bright sulphur colour. Only the young plants should be used for food.

Polyporus squamosus

Polyporus squamosus is found often on decayed ash trees and sometimes on others. The cap has a thick lateral dark stem and is pale yellow tinged with brown, and covered with dark scales. It is reported to have been found seven feet in circumference, with a weight of forty-two pounds, and that it attained this growth in the short time of four weeks.

Polyporus lucidus

Polyporus lucidus has a lateral stem which, with all but the margin of the cap, is highly polished, as if varnished, and is in colour a rich mahogany brown.

Polyporus arcularius (See Plate Facing Page 142)

Cap—Dark brown, minutely scaly, depressed in the centre; margin stiff, edge hairy, no flesh.

Tubes—Dingy cream colour. Openings oblong, almost diamond-shape, resembling the meshes of a net, drawn from stem to the margin of the cap, the meshes smaller on the margin, and simply marked out at the top of the stem.

Stem—Dark brown, minutely scaly, mottled, with a ground work of cream colour. Older stems are roughened at the base with whitish hairs. Hollow.

Spores—Creamy white.

Habitat—The specimen pictured was found growing on decayed branches of an oak tree in North Carolina.

Polyporus versicolor*

Polyporus versicolor has a leathery cap, thin and rigid, plane, depressed at the attached portion, velvety, shining with variegated two-coloured zones. The pores are minute, round, with acute and ragged edges. White, then yellowish. It is common on decaying tree trunks and upon telegraph poles.

Squā-mō'-sŭs Lū'-çĭ-dŭs Ăr-cū-lā'-rĭ-ŭs Vĕr-sĭc'-ŏ-lŏr

* Known also as Polystictus versicolor



Polyporus versicolor, Fr.



Polyporus circinatus, Fr.

See page 113



Lenzites betulina, Fr.

See page 113

Polyporus circinatus (See Plate Facing Page 112)

Polyporus circinatus quite often has one cap within another. The caps are thick, round, without zones, velvety, and of a rusty-yellow colour. The lower surfaces of the caps, with the stems are woody and corky, but the upper surfaces of the caps are soft and woolly. The stem is swollen and covered with yellow wool. The pores are decurrent, minute, and unbroken, and of a dusky grey colour. This is a handsome species, especially remarkable for its double cap. The stem is an inch long and almost equally thick. The cap is three to four inches wide, and the flesh is uniform in colour.

GENUS TRAMETES

Trametes pini is brown, and grows on pines and other cone-bearing trees. *Trametes cinnabarina* is bright red, and common on birch and cherry. *Trametes suaveolens* is white, and grows on willows.

GENUS LENZITES

Lenzites betulina

The *Lenzites betulina* has a somewhat corky, leathery cap, firm, and without zones, woolly and pale; the margin of uniform colour; the lamellæ radial, somewhat branching, and coming together again. Sordid white. This fungus, in the form of brackets, is found especially on birch trees, but it is also found on various other kinds. It is a widespread species, and is quite variable. The lamellæ are at first thick, corky, and sordid white; then thinner, with acute edges.



Under surface of L. betulina

Çĭr-çĭ-nā'-tŭs Pĭ'-nī Cĭn'-nă-bă-rī'-nă Sū-ā-vē'-ŏ-lĕns Lĕn-zī'-tēs Bĕt-ŭ-lī'-nă

Lenzites sepiria

Leathery shells, with the upper surface marked with rough zones of various shades of brown: the under surface with brown, radiating, papery gills.

GENUS DÆDALEA

The genus *Dædalea* has the spore-bearing surface in the form of winding and labyrinthine lamellæ, so that instead of pores there are irregular branching slits on the under surface. These fungi are normally sessile, woody, and hard.

Dædalea unicolor

Dædalea unicolor is in form like a full leather ruffle with scalloped edge, so full that the scallops overlap. The branching slits are very minute, and the upper side is in zones of tan colour, with a plush-like surface. The zones are often tinged with blue or green from algæ which grow upon them.


Dædalea confragosa

Dædalea confragosa has a rough grey upper surface, and grows from the centre in the form of depressed sessile caps. The texture is firm and leathery.

Dædalea quercina

Dædalea quercina is a woody or corky species, rugged, and without zones on the surface. At first it is porous; then, by the breaking down of the walls of the pores, slits are formed, with blunt partitions. It is common on oak trees.

Sē-pī'-rĭ-ă Dæ-dā'-lē-ă Cŏn'-fră-gō'-să Quĕr-çī'-nă



Dædalea quercina, (L.) Pers.

Upper side. Reduced



Dædalea quercina, (L.) Pers.

Lower side. Reduced

GENUS FAVOLUS

The genus *Favolus* has but few species ; one is very common on beech and hickory trees. *Favolus areolarius* has a lateral stem. The cap is depressed, and has a smooth, creamy-white upper surface, with long hexagonal pores running down the short stem.



Fă'-vŏ-lŭs F. alveolarius

CHAPTER IX: GELATINOUS AND OTHER FUNGI

Jew's Ear, or Judas's Ear (Edible)

Hirneola auricula-Judæ

THE Jew's ear is a gelatinous fungus which is so lobed and folded as to resemble a human ear. It is this resemblance to an ear which has suggested the names, Jew's ear and Judas's ear. Its habit of growing on elder has given rise to the belief that Judas Iscariot hanged himself on this tree. This fungus is prized by the Chinese as an article of food, and is imported by them from the South Sea Islands.

Its manner of bearing the spores on basidia places it in Class III, *Basidiomycetes*, which contains fungi with gills, teeth, and pores.


Tremellodon

Tremellodon is a clear fungus, which in colour resembles a piece of water-soaked snow, but in form resembles a *Hydnum*, as it has teeth-like projections on the under surface.

Guepinia


Guepina is a bright yellow fungus, about an inch high, in form like little spatules or goblets. It is common in cracks of logs and stumps.

Hĭr-nē'-ŏ-lă Au-rĭc'-ŭ-lă Trē-mĕl'-lō-dŏn Guĕp-ĭn'-ĭ-ă



JEW'S EAR. JUDAS'S EAR (EDIBLE)

(*Hirneola Auricula-Judæ*, (L.) Berk.)



XYLARIA

See page 136

CHAPTER X: OFFENSIVE FUNGI—ORDER PHALLALES

THE *Phallales* are all terrestrial fungi; that is, they are found growing on the ground, and not on logs and trees. They are interesting, but too offensive to attract any but the most courageous students.

They are more common in the South than in the North, where they make themselves too conspicuous by their intolerable odour.

They are structurally different from the other pouch-fungi, the puffballs, in that they do not retain their spores in the skin (peridium) until they are fully mature, but send them into the light, exposed on a quickly deliquescing jelly-like mass, the gleba.

The order may conveniently be divided into two families:

FAMILY I. CLATHRACEÆ { Receptacle latticed, sessile, or stalked. Spore mass borne on the inner surface of the receptacle.

FAMILY II. PHALLACEÆ { Receptacle tubular or cylindric, with a cap. Spore mass on the surface of the cap.

GENUS PHALLUS

Stinkhorns

The genus *Phallus* may be readily distinguished by the cylindrical shape of the spore receptacles and the intolerable odour. No one with his sense of smell developed would think of eating the members of this group. The botanist and the artist, however, have braved this lion on his own territory, and have found much that is beautiful and interesting; the artist having the advantage in his task of portraying the handsome specimen,

Phăl-lā'-lēs Clăth-rā'-çĕ-æ Phăl-lā'-çĕ-æ Phăl-lŭs

in that he can inclose it in a sealed glass case and work in comfort. The experience of the botanist must be realised to be appreciated.

An overpowering fetid odour suddenly evident upon the premises has many times filled with consternation the guests at summer resorts, causing among them much speculation, with suggestions of bad sewerage, and carelessness on the part of their host, together with other comments equally disastrous to the reputation of the place.

The distracted householder searches in vain for a solution of the difficulty, and the odour disappears as mysteriously as it came. If he is one of the initiated, however, he will search until he finds the haunt of the offender, and will destroy all chance of a repetition of the nuisance—for one summer, at least.

The mischief-maker is a handsome specimen, as its plate shows. The white stem, bearing at its summit a mass of gelatinous green substance capped with a yellow-white ring, and emitting its intolerable odour, has surely come into existence for a purpose—a purpose soon suggested—as hundreds of flies wing their way hither to sip the semi-fluid mass.

The botanist tells us that the spores of this plant are mixed in the green fluid, and that they are carried away on the feet and in the bodies of the flies to other places, where new colonies may be started.

The plant has undoubtedly emerged from the ground for the sole purpose of disseminating its spores, and all its parts have been developed to accomplish this function in the most effectual manner.


The banquet for the flies is prepared underground, and the table, with its viands all ready, is pushed into the light, while the invitation to the guests is wafted swiftly on the breeze.

One is curious to learn the mechanism by which so much is accomplished in apparently so short a time, and finds in this instance, as in all others where great things are accomplished with ease, that many forces have been slowly at work to insure everything being in readiness for the success of a final flourish. A search underground shows the mycelial threads to have permeated the soil for many feet in every direction in search of building material, and a glance at a vertical section of one of the pink eggs which has pushed its way out of the soil will show



Phallus impudicus, L.


in outline the plan of what is to be. The pink "eggshell," or peridium, is lined with a jelly-like substance, which has undoubtedly served as a safe packing to what is within, a protection against blows and insect ravages. Within this coat, in section, appears as two dark-coloured saddle-bags that which later is to form the green mass on the cap of the fully developed spore table. Between these dark masses lies in section the future stem; it is hollow, and bears on its rim the spore-bearing cap. The walls of this cap consist of flattened cells, which by extraordinary growth and expansion are to force the stem through the eggshell and carry the banquet of spores several inches into the light, leaving the torn wrapper as a volva at its base, a natural "Jack-in-the-box."



Section of young phallus

Phallus impudicus

Cap or Pileus—Outer surface bearing the spores in a jelly-like mass, gleba. Conic-campanulate. Outer surface sculptured with reticulated ridges after the green spore mass has disappeared.
Stem—Hollow, tapering at each end; upper end joined with the cap by a recurved border.
Veil—Wanting.
Volva—Pinkish.
Habitat—Low ground.



P. impudicus

Ĭm-pūd'-ĭ-cŭs

GENUS DICTYOPHORA

The genus *Dictyophora* differs from the genus *Phallus* in having a veil suspended from the apex of the stem, underneath the pileus or cap.

Dictyophora Ravenelii

Stem—Tapering at both ends.
Veil—Short, not reaching below the cap.


Dictyophora duplicata

Veil—Voluminous, hanging for several inches below the cap.


It is thought that this delicate white network, which hangs like a lace skirt below the cap, renders the fungus additionally conspicuous after dark, thus attracting the night moths and other night-flying insects.

Mutinus Caninus (See Plate Facing Page 136)

Cap or Pileus—Flesh coloured. The spore-bearing mass, oblong, ovoid, or conical, occupying one-third to one-sixth the total length of the stipe.



M. caninus



M. bambusinus

Stem or Stipe—Hollow, cylindrical, fusiform.
Habitat—About buildings, in gardens and thickets.

Dĭc-tў-ŏph'-ŏ-ră	Dū-plĭ-cā'-tă	Cā-nī'-nŭs
Ră'-vĕn-ĕl'-ii	Mū-tī'-nŭs	Băm-bū-sī'-nŭs


FAMILY CLATHRACEÆ

The members of the family *Clathraceæ* have a volva similar to the volva of the *Phallaceæ*. The volva ruptures, and the receptacle issues in a similar manner. The members of this family have the spore receptacle latticed or branched instead of tubular or cylindrical, and bear the spores on the inner surface of the receptacle rather than on the outer surface.

Latticed Clathrus

Clathrus cancellatus


Receptacle not stalked. The bars of the lattice-work are oblique and transversely wrinkled. The outer surface may be cinnabar red or white or yellowish. The inner surface of the bars is red.



C. cancellatus

Clathrus columnatus

Receptacle not stalked, consisting of from 2–5 vertical columns, separate below, but jointed at at the apex. Columns cinnabar red, enclosing the spore mass. Odour very fetid.




A. borealis

Anthurus borealis

Receptacle stalked, hollow, divided above into arms, which do not join at their apices, and which bear the spore masses on their inner surfaces, enclosing the spore mass when young, but later diverging.

Stem of receptacle white; arms narrow lance-shaped, with pale flesh-coloured backs, traversed their entire length by a shallow furrow.

Clăth-rā'-çĕ-æ	Căn-çĕl-lā'-tŭs	Ăn-thū'-rŭs
Clăth'-rŭs	Cŏl-ŭm-nā'-tŭs	Bō'-rĕ-ā'-lĭs



S. rubescens

Simblum rubescens

Receptacle stalked and globular, bars of the lattice forming meshes of about equal diameter either way. Red or flesh coloured, and transversely wrinkled.

Sĭm'-blŭm Rū-bĕs'-çĕns

CHAPTER XI: PUFFBALLS

ORDER LYCOPERDALES

THE pouch fungi include all fungi which have their spores or seeds in closed chambers until maturity—that is, until they are fully ripe and ready to be scattered by winds or animals. Collectively, the closed chambers are called the gleba, and this gleba is surrounded by a definite rind (peridium), which, in different puffballs, has various and characteristic ways of opening to permit the spores to escape.

The different ways in which the rinds (peridia) open are explained under the separate examples of the pouch fungi—puffballs, earth-stars, stinkhorns, birds' nests, and calostomas.

The *Lycoperdales*, known in different parts of the country as smokeballs, devil's snuffboxes, puffballs, etc., have their spores enclosed until maturity in closed chambers, surrounded by a continuous skin or peridium. They spend most of their lifetime underground, getting their food from decaying vegetable matter, and are for this reason called subterranean saprophytes. When they are about ready to scatter their spores, they emerge from the ground, and are then to be seen in pastures, and on fallen logs in woods and along roadsides. Every country child has pinched them to see the "smoke" rise, little knowing that he was doing for the puffball just that for which it had come into existence—scattering its spores far and wide to grow into new plants.

The plants of the puffballs, the mycelial threads, form an extensive network of white threads in the decaying vegetable matter in which they grow; then little balls appear on the white threads, as in the *Agaricales*, with the difference that they increase in size without forming gills and stem. The balls have a fleshy interior, cheesy and white at first, but afterwards yellowish or pinkish, gradually darkening until the whole or a part

Lȳ'-cō-pĕr-dā'-les

of the fleshy interior becomes filled with dust-like spores, when the rind of the ball breaks to let the spores escape.

Sometimes the wall breaks off in scales; sometimes it is punctured at the summit with one hole, sometimes with several, and sometimes it splits and turns back to form a star on the ground. Sometimes the balls contain elastic threads (capillitium), which help to push out the spores, and sometimes they do not. Sometimes there are threads massed at the base without spores in them, so that they form a sterile base or sterile subgleba, and sometimes the threads are massed to form a central column (columella) in the interior of the ball. These characters, with others, form the basis on which the puffballs are separated into the genera *Lycoperdon*, *Geaster*, *Calvatia*, *Bovistella*, *Bovista*, and *Calostoma*.

GENUS LYCOPERDON


The Lycoperdons, or true puffballs, produce within the ball vast numbers of dust-like spores mingled with elastic threads. When the ball is compressed, the rind or peridium bursts at the summit to form a single mouth, and the elastic threads cause the spores to fly out in puffs like smoke.



Lycoperdon


The spore-bearing part of the plant is globe-shaped, obovoid, or top-shaped, and at the base of the gleba no spores are produced; the cells here are coarse and empty.

The rind or peridium of the ball consists of two parts, the outer bark or outer peridium being adorned with spines or scales or warts or granules. Sometimes the exterior coat may be peeled off, sometimes it dries and falls away in fragments. The inner coat is thin and papery, and opens on the top with one opening. At first the ball is fleshy within, the microscope showing the flesh to consist of a great number of simple or branched threads and enlarged cells. The enlarged cells bear usually four



Section of Lycoperdon (diagrammatic)

Lȳ'-cō-pĕr'-dŏn



PEAR-SHAPED PUFFBALL (EDIBLE)

(*Lycoperdon pyriforme*, Schaeff. Nat. size)

spicules, on the tips of which are the spores. When the plant is fully developed, the fleshy part becomes so filled with moisture that water may be squeezed out as from a sponge. As the flesh becomes moist, the colour changes from white, through yellow, to olive. After the change in colour, the wet mass becomes dry and powdery, a mass of globose spores and elastic threads or capillitium.

The Lycoperdons are of small size, usually found in fields and woods. A section made by cutting a ball from top to base will show that the threads form a more or less well developed sterile base or subgleba underneath the fertile gleba, or mass of threads containing spores. Sometimes the sterile threads from the base rise upward in the centre of the fertile mass and form a little column, the columella. Usually the threads which bear the spores are in two sets ; one set extending inward from the walls of the rind, and another set extending outward from the central columella.

Pear-shaped Puffball (Edible)

Lycoperdon pyriforme

Peridium or Pouch—Pear-shaped. Dingy white or brownish, with mycelium of long, white, branching fibres. Diameter ¾–1¼ inches. Height 1–2 inches.
Bark or Outer Coat—Thin; of minute, often persistent, scales or granules, or short, stout spinules. Whitish grey or brownish.
Inner Coat—Smooth, papery, whitish grey or brownish, opening by apical mouth.
Subgleba—Small, white, quite compact, the cells minute.
Columella—Present.
Spores—Globose, even, greenish yellow to brownish olive.
Threads—Branched, long, forming a dense tuft in the centre.
Time—July to October.
Habitat—On old timber or on the ground, in groups sometimes several feet across in extent. The commonest of puffballs, and found throughout the world.

Pinkish Puffball (Edible)

Lycoperdon subincarnatum (See Plate Facing Page 134)

Peridium or Pouch—Globe-shaped, sessile, without a stem-like base. Rarely over one inch in diameter.


Pȳ'-rĭ-fŏr'-mĕ Sŭb-ĭn'-căr-nā'-tŭm

Bark or Outer Coat—Pinkish brown, with minute short, stout spinules, which fall away at maturity.
Inner Coat—Ash coloured. Deeply pitted by the falling off of the spinules of the outer coat, the pits not surrounded with dotted lines.
Columella—Present.
Subgleba—Small but distinct.
Spores and Capillitium—Spores round, minutely warted. Greenish yellow, then brownish olive. Threads long, simple, and transparent.
Time—August to October.
Habitat—Old trunks in woods.

GENUS CALVATIA

The *Calvatias* are puffballs of large size, all with thick cord-like mycelium rooting from the base. They all eject their spores through irregular openings in the upper part of the peridium, and they all have a dense network of branching threads (capillitium) traversing the tissues of the spore-bearing portion, the gleba. These threads are elastic, and project the spores from the rind or case as they twist and turn. The sterile portion, the subgleba, is definitely limited and concave above.


Calvatia




Section of Calvatia

Brain-shaped Calvatia (Edible)

Calvatia craniformis

Peridium or Pouch—Very large, obovoid or top-shaped, depressed above.
Bark or Outer Coat—Smooth, very thin and fragile, easily peeling off. Pallid or greyish, often with a reddish tinge ; often

Căl-vā'-shĭ-ă Crā-nĭ-fôr'-mĭs



BRAIN PUFFBALL (EDIBLE)

(*Calvatia craniformis*, Schw.)

wrinkling to resemble somewhat the surface of the brain, whence its name *craniformis.*

Inner Coat—Thin, ochreous to bright brown, velvety, extremely fragile. The upper part breaks into fragments.

Subgleba—Occupies half the peridium ; cup-shaped above, persistent.

Spores—Greenish yellow, then olivaceous. Globose, even, with minute pedicel.

Threads—Long.

Habitat—On ground in woods.

Giant Puffball (Edible)

Calvatia maxima

Peridium or Pouch—Very large, 8–15 inches in diameter, or larger. Globose, depressed globose, or obovoid, with a thick cord-like root.

Bark or Outer Coat—Flocculous or nearly smooth, thin, and fragile. White or greyish, becoming yellowish or brown; usually remaining closely adherent to the inner coat.

Inner Coat—Thin and fragile after maturity, breaking up into fragments.

Subgleba—Shallow or none.

Spores and Threads—Greenish yellow, then brownish olive. Spores globose, threads long and branched.

Time—August to September.

Habitat—Grassy places.

Calvatia maxima has been known as *Lycoperdon giganteum,* and also as *Lycoperdon maxima.* It has been transferred from the genus *Lycoperdon* to the genus *Calvatia* because it ruptures the peridium irregularly to discharge its spores, instead of forming a small hole at the apex, as other Lycoperdons do.

It is asserted on good authority that the giant puffball has been found with a diameter of three feet and a weight of forty-seven pounds. The giant puffball is considered by many as a choice article of food when the flesh is white. It is said that if the flesh of a growing puffball is cut or injured the wounds will fill up with new tissue. It will be interesting for some one to try this experiment.

In the days before matches came into use, the dry, spongy threads were used as tinder to catch the sparks which flew from the flint-stone when it was struck for fire, and the spore-dust was used to stanch the flow of blood.

Cup-shaped Puffball (Edible)

Calvatia cyathiformis


Peridium—Large, top-shaped.
Bark or Outer Coat—Thin, adherent, smooth, and continuous, easily peeling off.
Inner Coat—Pale to dark purple, loosely woven, fragile at maturity, breaking up into fragments from above downward.
Subgleba—Short and thick, with cord-like root, persistent, cup-shaped, occupying ⅓–½ the peridium.
Spores and Threads—Violet to dark purple. Spores globose and warted, threads long.
Time—August to October.
Habitat—On the ground in meadows and pastures.

The old name was *Lycoperdon cyathiforme*. *Cyathiforme*, meaning cup-shaped, is suggested by the cup-like base which remains after the dispersion of the spores and threads (capillitium).

GENUS BOVISTA

In the genus *Bovista* the rind or peridium opens by an apical mouth, as do the species of *Lycoperdon*, but the species of *Bovista* have no sterile base. They are puffballs of small size, growing in fields and woods. The outer coat is thin and fragile, and scales off at maturity. The inner coat is thin, becoming papery, and then opens by an apical mouth.


Bovista



Section of Bovista (diagrammatic)

Bovista plumbea is esteemed a delicacy. It is shaped like a flattened globe, with a smooth, white inner coat, and a lead-coloured outer coat opening by an apical mouth.

Çÿ'-ăth-ĭ-fôr'-mĭs Bō-vĭs'-tă Plŭm'-bĕ-ă



Bovistella Ohiensis, Ellis and Morgan
(Edible.) Reduced



CUP-SHAPED CALVATIA (EDIBLE)
(*Calvatia cyathiformis*, Bosc)
Reduced. Nat. size: Ball diam., 2⅞ inches

GENUS BOVISTELLA

The genus *Bovistella* contains but one species.


Bovistella Ohiensis (Edible) (See Plate Facing Page 128)

Peridium or Pouch—Globose or broadly obovoid, sometimes much depressed, wrinkled underneath, with thick cord-like base.

Bark or Outer Coat—Dense, floccose, or with soft warts or spines. White or greyish, drying to buff colour, and falling away.

Inner Coat—Smooth, shining, pale brown or yellowish surface.

Subgleba—Cup-shaped, broad, ample, occupying nearly one-half the peridium; long, persistent.



Section of Bovistella (diagrammatic)

Spores and Capillitium—Loose, friable, clay colour. Threads free, short, twice branching, originating within the spore mass, and having no connection with the tissue of the inner coat.

Habitat—On ground in pastures and open woods.

GENUS GEASTER

Earth-stars

The *Geasters* or Earth-stars are the most picturesque forms of the puffballs. At first they are sunk deep in the soil, and are connected with it by abundant thread-like mycelium, which issues from every part of the surface. In the earth-stars the covering to the pouch is double, the outer cover is thick and leathery, and at first closely invests the inner coat, but is separate from it. At maturity the outer coat breaks its connection with the mycelium in the soil and bursts to form separate lobes, which become reflexed and lift the inner ball from the ground into the air, where it remains, seated at the centre of the expanded star-

Bō-vĭs-tĕl′-lă Ō-hī-ĕn′-sĭs Ġē-ăs′-tĕr

like coat. The coat of this ball is thin and papery, and opens by one apical mouth. The threads or capillitium, which bear the spores, project from the tissue of the inner wall and also from a central columella.

The *Geasters* have no economic value, but are rather interesting to the nature student on account of their beauty and their curious methods of discharging their spores.

The Smallest Earth-star

Geaster minimus

Peridium or Pouch—Globose, depressed, not pointed; vaulted underneath.
Bark or Outer Coat—Segments acute at the apex, many lobed; the lobes, seven to nine, expanding ½–1 inch.
Inner Coat—Ovoid, ¼–1 inch in diameter, white to pale brown, with a distinct pedicel, seated in a plain circular disk.
Mouth—Lifted on a cone, lip bordered with hair-like fringe.
Spores—Brown, globe-shaped, minutely warted.
Threads—Transparent.
Habitat—Grassy grounds.

Water-measuring Earth-star

Geaster hygrometricus

Peridium or Pouch—Sub-globose, depressed, the bark or skin falling with the mycelium.
Bark or Outer Coat—Deeply parted; the segments, acute at the apex, seven to twenty. Strongly hygrometric, expanding to a breadth of 2–3 inches.
Inner Coat—Globose, depressed, sessile, covered with a network. Whitish or greyish.
Mouth—Rim irregular.
Spores—Brown, globe-shaped, minutely warted.
Threads—Transparent, much branched and interwoven; continuous with the hyphæ or threads of the inner coat.
Habitat—Fields and woods, in sandy soil.


The *Geaster hygrometricus*, or *Astræus hygrometricus* as it is called by some, is found all over the world. When the

Mĭn'-ĭ-mŭs Hȳ'-grŏ-mĕt'-rĭ-cŭs




LEAST EARTH-STAR

(*Geaster minimus*, Schw. Nat. size)



WATER-MEASURING EARTH-STAR

(*Geaster hygrometricus*, Pers. Nat. size)



BIRD'S NEST

(*Cyathus vernicosus*, D. C. Nat. size)

See page 133

weather is wet, the lining of the points of the star become gelatinous and lie flat on the ground, anchoring the plant firmly ; but when the weather is dry, the soft, gelatinous part becomes hard and rigid, and curls the segments up around the inner ball; then the wind rolls it about, and it scatters its spores from the hole in the apex of the ball as it rolls. It is a fair-weather traveller, always resting at night and on damp days.

GENUS CALOSTOMA

This genus has but three known American species. The plants are remarkable in structure and substance. The spore mass or gleba lies at the centre of a base, and is in its young stages surrounded by four layers. The outermost coat is gelatinous, and soon disappears. This is known as the volva or wrapper. The layer just within the wrapper also soon disappears ; it is known as the exoperidium. The layer just within the exoperidium is known as the endoperidium, and is the layer seen on the exterior of older specimens. Between the exoperidium and endoperidium is a layer of red threads, part of which is torn away when the exoperidium breaks, and a part of which remains as a red star at the apex of the ball.

The exoperidium is cartilaginous ; it is thin and fragile when dry, but when wet it is flexible, translucent, and soft. The endoperidium is hard and rigid when dry, and is conspicuous for the brilliant red which shows at its mouth. Within the endoperidium is a sac which contains the spores ; when the spores are mature, this sac contracts, and forces the spores out into the air. The mycelium, or vegetative part of the *Calostoma* plant, is composed of numerous cord-like fibres, translucent, jelly-like, and tough, which, branching and anastomosing into a dense network, form a rooting columnar base to the spore-bearing part of the plant. The name *Calostoma* means beautiful mouth, referring to the red star which surrounds the opening through which the spores escape.

Căl-ŏs'-tŏ-mă

Calostoma lutescens

Ball or Peridium—Globular, with a thick, entangled, rooting base.
Outer Coat or Exoperidium—Dingy yellow, rupturing so as to form a ragged collar at the base, a thin cap on the summit, and small fragments between. The cap is marked on the under side with a vermilion star outlined with yellow.
Inner Coat or Endoperidium—Smooth, globular. Pale yellow.
Mouth—With several rays of a bright red colour.
Footstalk—Thick, of entangled strands finer than those of *C. cinnabarinum*. Stem long, yellowish green.
Spores—Globular, with protuberances.


Calostoma Ravenelii

Ball or Peridium—Globular, with a thick, entangled, rooting base.
Outer Coat or Exoperidium—Cartilaginous, gelatinous. *Fragments remain upon the inner coat* in the form of irregular warts or scales.
Inner Coat or Endoperidium—Yellowish. Tough when wet, rigid when dry.
Mouth—Red starred.
Footstalk—Short and rooting, composed of mycelial threads netted to form cartilaginous cords.
Spores—Elliptical, oblong, smooth.

Calostoma cinnabarinum

Outer Coat or Exoperidium—Vermilion within, breaking at the base and sometimes at the apex.
Inner Coat or Endoperidium—Yellowish, often slightly vermilion. *Smooth.*
Mouth—Rays several. Vermilion.
Footstalk—Reddish brown or brownish.
Spores—Elliptical oblong, spined or punctured, pale ochre yellow.

Lŭ-tĕs'-çĕns Răv'-vĕn-ĕl'-ī-ī Cĭn'-nă-bă-rī'-nŭm



Calostoma Ravenelii (Berk.) Mass.



Calostoma lutescens (Schw.), Burnap.



Calostoma cinnabarinum, Desv.



Spathularia velutipes, C. & F.

Reduced. See page 138.

ORDER NIDULARIALES

The members of this order, *Nidulariales*, or bird's-nest fungi, are curious fungi of small size. They resemble, when mature, tiny birds' nests containing eggs, as the pouch in which the spores are developed opens at the top to form a nest or bowl or trumpet, and the globular cases in which the spores are contained have strong walls, and remain in the open pouch like eggs in a nest.

One genus of the order, *Sphærobolus*, has two walls or layers to its nest and but one spore case or "egg." This "egg" is jelly-like, and is forcibly thrown from the nest when the spores in it are mature. While a part of this action is due to the mechanical working of the teeth, it is thought that underneath the spore case gases are formed which expand, and so help force out the "egg."

The other genera have but one wall to their nests. The genus *Nidularia* has ragged edges, the genus *Cyathus* is trumpet-shaped, and the genus *Crucibulum* is bowl-shaped.

ORDER SCLERODERMATALES

The puffballs of the order *Sclerodermatales* have the rind or peridium thick. The spores remain in the peridium until maturity, when they escape from an irregular opening in the rind. The species are not numerous, but some are abundant and widely distributed.

GENUS SCLERODERMA

The species of the genus *Scleroderma*, or thick-skinned puffballs, represent a transition from the subterranean forms to those which emerge from the ground and have a definite opening from which they eject their spores. To illustrate: there is one group (*Hymenogastrales*) in which the species remain in the ground,

Nĭd-ū-lă'-rĭ-ă'-lēs Nĭd-ū-lă'-rĭ-ă Crū-çĭb'-ŭ-lŭm Scl̆ĕr-ŏ-dĕr'-mă
Sphē-rŏb'-ŏ-lŭs Çȳ'-ă-thŭs Scl̆ĕr'-ŏ-dĕr'-mă-tă'-lēs

even when mature, unless washed out by rains or nosed out by animals, and which rely upon being crushed by accident or upon the disintegration of their rinds to free their spores. All true puffballs push themselves out of the ground before the spores are mature, and then, by rupturing the skin in a definite manner when mature, disperse their spores. The thick-skinned puffball goes a step farther than the first mentioned and emerges from the ground; but it remains behind the true puffballs, which disperse their spores from a definite opening, for it ruptures the skin irregularly.

Scleroderma vulgare (Edible)



Section of S. vulgare

Peridium or Pouch—Rough and warty; depressed, globose. Pinkish to buff, remaining solid until the fungus is quite old. Sessile or with a rooting base. Ruptures irregularly to scatter the spores.
Subgleba—None.
Spore Mass—Lead colour marbled with white.
Habitat—Old stumps and buried roots.

This fungus has been eaten without harm, but is pronounced very unattractive.


Vŭl-gā'-rē



FLESH-COLOURED PUFFBALL (EDIBLE)
(*Lycoperdon subincarnatum*, Peck)
See page 125




YOUNG PEAR-SHAPED PUFFBALL (EDIBLE)
(*Lycoperdon pyriforme*, Schaeff)
See page 125



HARD-SKINNED PUFFBALL
(*Scleroderma vulgare*, Fries)

CHAPTER XII: SPORE-SAC FUNGI—ASCOMYCETES


ALL the fungi which belong in the class *Ascomycetes* develop their spores in little membranous sacs called asci. These asci are, as a rule, collected, together with slender empty asci, called paraphyses, in variously shaped bodies, known in different orders by different names, such as perithecium, ascoma, apothecium, and receptacle. This class includes in its numbers individuals ranging in size from microscopic one-celled plants to conspicuous and often beautiful specimens. We shall concern ourselves with but a few of those conspicuous ones which are attractive either from an edible or an artistic standpoint.




Asci and paraphyses

ORDER TUBERALES—TRUFFLES

The order *Tuberales* contains the truffles, which are subterranean fungi, ranging in size from an acorn to a good-sized potato. The asci or spore-sacs are formed on the interior of the fungus, the warty truffle itself being called an ascoma, as it contains the asci.


Section to show position of asci


Truffles, ascoma


Since the time of Pliny and Dioscorides, truffles have been known and esteemed as a table delicacy. Since they mature

Ăs'-cŏ-mȳ-cē'-tēs Tū'-bĕr-ā'-lēs

underground, they must be hunted for by dogs and pigs trained for the purpose. A pig will scent a truffle at a distance of twenty feet, and will run quickly to the spot to dig it out with her snout. An attendant must follow the pig to secure the truffle before the pig eats it. Edible species have not been found growing in this country.

ORDER HYPOCREALES

The order *Hypocreales* contains certain fungi which are parasitic on other fungi, and also on insects. In the genus *Cordyceps* there is a club-like form about an inch long, and of a rich red colour, which lives as a parasite on the pupæ of various moths buried beneath dead leaves. In New Zealand a similar fungus lives on caterpillars. The mycelium in time replaces the body, but maintains outwardly the form and appearance of the caterpillar. These growths are much prized by the natives as food. The form which grows on the truffle-like *Elaphomyces* is shown in the species of *Elaphomyces* upon which the *Cordyceps* lives as a parasite grow two or three inches below the surface of the ground, and somewhat resembles a truffle in appearance.



Fungus growing on caterpillar

ORDER SPHÆRIALES

(See Plate Facing Page 116)

The *Xylaria* pictured is a woody fungus which is common, growing on logs or at the bases of trees or stumps. The collections of asci (*perithecia*) are fully imbedded in the fleshy part of the fungus (*stroma*), which is formed almost wholly of hardened mycelium.

Hȳ-pŏ-crĕ-ā′-lēs Cŏr′-dȳ-çĕps Ē′-läph-ō′-mȳ-cēs Sphē-rĭ-ā′-lēs Zȳ-lā′-rĭ-ȧ



Cordyceps capitala (Holmsk.), Lk.
(Parasitic on Elaphomyces.)
Reduced.



Lycogala epidendron.
Reduced. See page 144.



Floccose Chanterelle (edible).
(Cantharellus floccosus, Schw.)
See page 53.



Mutinus caninus, Huds.
See page 120.



SLIPPERY LEOTIA (EDIBLE)

(*Leotia lubrica* (Scop.), Pers.)

Gelatinous, gristly; spore-bearing body (*ascoma*), green or yellow; stem yellow
Family, *Geoglossaceæ*. Class, *Ascomycetes*. Order, *Helvellales*. See p. 138



JELLY-LIKE TREMELLODON (EDIBLE, Mcl.)

(*Tremellodon gelatinosum*)

Surface white to grey; teeth white. Class, *Basidiomycetes*. Order, *Tremellales*. See Genus, p. 116


ORDER PEZIZALES—CUP FUNGI

The *Peziżales* or cup fungi, which are typically disk or cup-shape, comprise an extensive group, and vary in size from forms scarcely visible to the naked eye to forms several inches in diameter. One genus, *Peziża*, has a smooth ascomata, regularly saucer-shaped or cup-shaped. The genus contains many species which are met with chiefly on decaying vegetable matter. The spore-sacs are situated on the upper surfaces of the cups, and the spores are ejected with such force and in such profusion that they form a cloud around the plant from which they are expelled.

Peziża æruginosa is a stalked green form, and is interesting because it permeates the wood of oaks and beeches with its mycelium, and gives a rich green colour to the wood, which makes it valuable for the manufacture of the famous "Tunbridge ware." One may often find the wood affected when the fruit cups are not evident. *Peziża Willkommii* produces on larch trees a disease known as the "larch canker," which shows itself as a sunken, blistered hole from which resin flows. This patch is formed because the mycelium of the fungus attacks and destroys the cambium or green layer which lies under the brown bark, and since this is the tissue which builds up the wood of the trees, the growth of wood in this part is prevented. The fruits appear above the bark in the form of little cups, white outside and scarlet within. When branches bearing golden-yellow needles are seen among the fresh green shoots of a larch, prematurely giving to the tree an autumnal appearance, one may expect to find *Peziża Willkommii* at work.

Peziza odorata (Edible)

Cup—Yellowish, translucent, becoming dull brown when old. The flesh is moist and watery, and separable into two layers; the outer layer rough, and the inner smooth. The frame is cup-shaped when young, but flattened and split on the margin when old.



Section to show two layers

Ē′-rū-ġĭn-ō′-să Wĭl-kŏm′-ĭ-ī Ō′-dŏr-ā′-tă

Stem—None.
Habitat—The specimen pictured was found in December, among the violets in a cold frame in New Jersey.

Golden Peziza (Edible)

Peziza aurantia

Cup—Orange red within, golden or whitish outside, with a frosted appearance. Subsessile, irregular.
Habitat—In clusters on the ground, usually in the autumn.

ORDER HELVELLALES—EARTH-TONGUES

The order *Helvellales* contains the fleshy spore-sac fungi which have the spore-bearing body, the ascoma, open from the earliest stage of its development.

FAMILY GEOGLOSSACEÆ

The family *Geoglossaceæ*, which belongs to this order, contains the earth-tongues, which are club-like forms, green or black or yellow, and from one to three inches high, common on the ground, growing in rich wood mould. In consistency they are fleshy, gelatinous, or waxy, and their asci open by means of a terminal pore.

GENUS SPATHULARIA

The genus *Spathularia* has the spore body flattened and growing down both sides of the stem.

Velvety Spathularia

Spathularia velutipes (See Plate Facing Page 132)

Spore Body—Flattened, tawny yellow, shaped like a spatula, with the spore surface wavy and growing down two sides of the stem.
Stem—Hollow, *minutely velvety*, dark brown tinged with yellow.
Flesh—Firm and tough, shrinking little in drying.
Habitat—Mossy trunks in damp woods.

Au-rĭn'-shĭ-ă Vĕ-lū'-tĭ-pēs



Golden Peziza (edible).
(Peziza aurantia, Pers.)



Peziza cdorata. Pk. (edible).
Reduced. See page 137.

Spathularia clavata

Spore Body—Clear yellow, shaped like a spatula, sometimes tinged with red. Obtuse or cleft at the apex, the surface wavy, the margin crisped, growing down the stem some distance on opposite sides; hollow.
Stem—Thick, hollow. White, often becoming yellowish.
Flesh—Dry.

GENUS GEOGLOSSUM

The genus *Geoglossum* has the spore body simple, erect, and club-shaped, and entirely black. The spore surface is terminal.

Geoglossum hirsutum

Geoglossum hirsutum is black, dry, and everywhere velvety. Lanceolate, ovate, oblong, or almost round, often irregular. The spore-bearing portion is one-quarter to one-half the length of the fungus.

Geoglossum glabrum

Geoglossum glabrum is dry, black, or brownish black, sometimes tinged with olive or purple. Club-shaped or sometimes laterally compressed. The spore-bearing portion not sharply distinct from the stem below.

GENUS VIBRISSEA

The genus *Vibrissea* contains fungi with vertical and simple stems, and horizontal caps with their thick margins rolled in toward the stem. The spore-sacs are borne on the upper surface.

Vibrissea truncorum

Vibrissea truncorum is a clear orange-red or sometimes yellow or brownish-red fungus, about an inch high, found on decayed wood, branches, or leaves which are submerged in water.

Clăv-ă'-tă Hĭr-sū'-tŭm Vĭ-brĭs'-sĕ-ă
Gĕ-ŏ-glŏs'-sŭm Glă'-brŭm Trŭn-cō'-rŭm

Vibrissea circinans

Vibrissea circinans is a pale yellowish flesh colour, or simply yellowish fleshy fungus found growing in circles or clusters, with convex caps and incurved, wavy margins, the concave under surface often minutely wrinkled. The stem is long, pallid, or reddish. The plant is found chiefly in pine woods.

GENUS MITRULA

The genus *Mitrula* has the spore body erect, black or bright coloured, and dry, spatulate, or cylindrical, often compressed laterally. The spore-bearing surface is sharply distinct from the scaly or mealy stem below.

Irregular Mitrula (Edible)

Mitrula vitellina, var. irregularis *

Spore Body—Bright egg yellow. Club-shaped, somewhat lobed, cylindrical or compressed; apex narrow, obtuse, smooth. No two plants are quite alike. Length, 1–2 inches.

Stem—Short, white, rather distinct, covered with fibres. Spongy and white within.

Habitat—In mossy places in woods during the autumn. The specimen pictured was found growing among fallen birch leaves, hemlock needles, and moss, in the dense woods at Lake Placid.

FAMILY HELVELLACEÆ—MORELS

A second family *Helvellaceæ* contains three important genera, *Morchella*, *Gyromitra*, and *Helvella*, in which are the largest and most highly prized spore-sac fungi known. They are distinguished from the earth-tongues by the cap-like form of the spore body or ascoma, but especially by the character of the spore-sac, which opens by a little lid instead of by a simple pore.

Çir-çĭ'-năns	Vĭ'-tĕl-lĭ'-nă	Ġў'-rō-mĭ'-tră
Mĭt'-rŭ-lă	Mŏr-kĕl'-lă	Hĕl-vĕl'-lă

* This species is also described under the names *Geoglossum irregulare* and *Geoglossum vitellinum*.



Helvel a elastica, Bull. **Reduced.** See page 142.



Helvella lacunosa, Holm. **Reduced** See page 143.



Mitrula vitellina, sacc., var. irregularis, Pk (edible). **Reduced.**

GENUS GYROMITRA

The genus *Gyromitra* contains seven species. These have the ascomata distinctly stalked, and the upper surface covered with gyrose folds. The largest spore-sac fungi belong in this genus.

Gyromitra esculenta (Edible)

Cap—Bay red, round, lobed, irregular, gyrose-wrinkled, attached to the stem in several places. Hollow, white, and uneven within.
Stem—Whitish, hollow, scurfy. Two or more inches long.
Flesh—Edible only when young and freshly gathered.
Habitat—In sandy soil, during the wet weather of May and June.



G. esculenta



M. deliciosa
Section to show margin of the cap united to the stem.

GENUS MORCHELLA

The genus *Morchella* has the cap covered with a network of blunt ridges enclosing irregular depressed spaces. The spore-sacs are developed in both ridges and depressions. All the species when young are of a buff yellow tinged with brown, but later they are darker. The stems are rather stout and hollow, white or whitish in some species, and attached to the cap at the apex only; but in others, attached to the rim as well. All the species



M. esculenta

are edible and highly esteemed. They must be looked for during wet weather, early in the season. They may be classed in two groups, according to the attachment of the stem.

I—MARGIN OF CAP UNITED TO THE STEM

Cap rounded or oval..... MORCHELLA ESCULENTA
Cap oblong or cylindrical..... MORCHELLA DELICIOSA
Cap conical or oblong conical ; broader than stem..... MORCHELLA CONICA
Cap conical or oblong conical ; scarcely broader than stem. MORCHELLA ANGUSTICEPS

II—MARGIN OF CAP FREE FROM THE STEM

Cap free from the stem to the middle.... ..... MORCHELLA SEMILIBERA
Cap free from the stem to the top..... MORCHELLA BISPORA



Helvella

GENUS HELVELLA

The genus *Helvella* contains twelve species. They all have lobed, irregular, or saddle-shaped caps, which are fleshy, and attached to the stem at the apex. They are contorted in such a way that no two of the species appear alike.

Helvella elastica

(SEE PLATE FACING PAGE 140)

Cap—Brownish grey. Cup-shaped, flattening out when mature; when young, the under surface is covered with little spines or hairs. Both surfaces are smooth when mature.

Stem—Slender, of the same colour as the cap, tapering toward the cap. Solid and white within.

Habitat—The specimen pictured was found growing beneath hemlocks and yellow birches, in Lake Placid forest, during September.

Dĕ-lĭsh-ĭ-ō'-să Ăn-gŭs'-tĭ-çĕps Bĭ'-spŏr-ă
Cŏn'-ĭ-că Sĕm'-ĭ-lĭb'-ĕr-ă Ē-lăs'-tĭc-ă



DELICIOUS MOREL (EDIBLE)

(*Morchella deliciosa*, Fr.)



Polyporus arcularius, (Batsch) Fries

See page 112


Helvella lacunosa (See Plate Facing Page 140)

Cap—Of one piece, thin and flexible like rubber cloth, folded to saddle the apex of the stem. The two saddle-flaps are attached on their margins at irregular intervals when young, and are puffed out like a balloon; but when mature, the pieces separate. Their outer surface is brownish grey, and their inner surface light grey, creased and folded.

Stem—Irregularly and deeply furrowed.

Flesh—Odour offensive.

Habitat—The specimen pictured was found growing imbedded in deep moss on old bark in the Lake Placid woods.



H. lacunosa (diagrammatic)

Lăc-ŭ-nō'-să

CHAPTER XIII: SLIME FUNGI—MYXOMYCETES

(See Plate Facing Page 136)

Whether the slime fungi are plants or animals is a question not yet decided. They are living organisms which have no chlorophyll, or leaf-green, and which in their vegetative state resemble certain groups of the *Protozoa*, or unicellular animals, which live in water. In their manner of reproduction they show resemblances to certain fungi, and the spore cases or sporangia of some resemble tiny puffballs in form and manner of ejecting their spores.

In the growing stage they consist of a naked mass of yellowish or whitish protoplasm, which creeps about in the dark, in accumulations of dead parts of plants, or under the bark of rotting stumps or logs. When a spore germinates, the membrane about it bursts, and a bit of naked and slimy protoplasm escapes. This tiny mass creeps about, absorbing food from its surroundings, and increasing in size until it may perhaps cover an area of many inches. After a time spores begin to form, then either the whole mass is transformed into a single spore case or a number of spore cases are formed. The spore cases of *Lycogala epidendron* are pretty things, resembling pink coral beads. When fresh, a case is filled with a thin pink paste; but when mature, with fine brown dust-like spores. When the spore case bursts, these spores escape, and if they fall in favourable places the life cycle begins anew.

Mўx'-ō-mў-çē'-tēs Prō-tō-zō'-ă Lў-cŏg'-ă-lă Ĕp-ĭ-dĕn'-drŏn



BRISTLY PANUS

(*Panus strigosus*, B. & C.; *Lentinus strigosus*)

Cap, gills, and stem creamy. See Genus, p. 67

CHAPTER XIV: FUNGI FOR THE HERBARIUM

THERE are no plants more difficult to preserve for an herbarium than the fleshy fungi, and yet my personal observation leads me to believe that there are many people who would be willing to undertake the task if they knew how to set about it; and there is no class of plants in which the assistance of the amateur may help the botanist more than in this, provided that, at the time of gathering a specimen, full descriptive notes are made of all the characteristics of the plant.

To aid one in quickly taking notes, it is well to have with one in the field some printed or written blanks. A convenient form is suggested by the following outline, which is the one used by the Boston Mycological Club:

Species

Collected by
No.
Locality
Date

COLLECTOR'S NOTES.

N.B.—When collecting, be sure to get the whole plant, base and all, uninjured; and to get young as well as mature specimens.

Note here at once the

Habitat. On { Tree (kind; dead or living). / Ground (kind of soil).

Place (wood, field, wet or dry, high land or low, etc.).

Under and near what trees?

Manner of Growth. { (Solitary, in clusters, troops, or cæspitose [growing from one root]).

Character. (Viscid, hygrophanous [transparent when moist], dry.)

Smell.

Taste.

Spores. Colour.

NOTE.—If the plant is perishable, sketch and describe it fully at once, and look

for indications of spore colour. After the plant is described it may be dried in hot air (over a stove for instance) and preserved or sent in for identification.

Sketch the plant, indicating markings of cap and stem. Draw or trace a vertical section through the centre of cap and stem, indicating thickness of flesh; shape and attachment of gills; nature of interior of stem; position of ring, volva, etc. Do this also for a young specimen (button), showing whether the margin of the pileus is straight or incurved.

N.B.—If not life size, note dimensions.

When the characters vary with age or with moisture, note the changes.

Pileus.

- Shape (flat, convex, concave, umbonate [raised in the centre], umbilicate [depressed in the centre], etc.).
- Is it viscid when moist and fresh; tough, fragile, fleshy, membranaceous; smooth, floccose, scaly, silky, fibrillose; even, rough, wrinkled, furrowed? etc.
- Is the margin entire, wavy, striate, inrolled, upturned, smooth, woolly, hairy, appendiculate? etc.
- Colour and markings.

Gills.

- Shape.
- Attachment (adnate, sinuate, decurrent, etc., or *free*).
- Are they distant or crowded, all of one length, branching or forked, connected by veins?
- Surface (smooth, powdery, marked in any way).
- Colour (young and old).
- Texture (thick, thin, brittle, etc.).
- Margin (entire, wavy, scalloped, toothed, fringed).
- With *Boleti* note colour, length, and size of tubes, shape and size of mouths, relation of pore surface to stem, etc.

Flesh.

- Consistency (firm, mealy, punky, etc.).
- Colour (in general; just under skin; near gills or tubes).
- Juice (taste and colour).

Stem.

- Texture (tough, flexuous, fragile, fleshy).
- Shape (tapering either way, straight or bent, swollen, etc.).
- Exterior (cartilaginous, fibrous or not, etc.).
- Colour and markings (striate, dotted, pruinose [with a bloom], fibrillose, or smooth, etc.).
- Interior (hollow, solid, stuffed, fistulose [tubular], etc.).
- Base (shape, markings, etc.).

Mycelium (thread-like, cottony, compact, root-like, sclerotioid [hard], coloured).

Veil. (Examine young specimens).

Ring. (Relative position, permanent or fugacious [quickly disappearing], etc.).

Volva. (Examine young buttons, base of stem, surface of pileus).

Remarks.

COLLECTOR'S OUTFIT.—For collecting fungi there is nothing better than a cheap splint basket with a cover. The size will depend upon the ambition and strength of the collector. In addition, a chisel for woody fungi and a trowel or broad-bladed knife will be found convenient; also a few small boxes for fragile species, and a package of thin, tough, uncoloured paper in which other specimens may be put. Sheets of six inches square and also of twelve by twenty-four are convenient sizes. Before the specimens are put in the basket, those of a kind should be compactly piled in the centre of a sheet, and the four corners of the sheet brought together and fastened by twisting them. The slip with the notes may either be put inside the package or fastened on the outside.

CARE OF SPECIMENS.—As soon as possible after reaching home, the packages should be taken from the basket and spread out in convenient places. If the specimens are to be used immediately for identification, begin with the most perishable, or they will be lost by decay. If they are to be preserved for future use, put them in the warmest place available where they will not burn. This may be under or over the kitchen stove, or in the furnace-room of the hotel or laundry; or, if any of these places is not practicable, then a drier may be improvised by placing over a lamp a frame made of wire screening.

COLLECTING SPORES.—At least one specimen of each kind should be set for spores. This is readily done by thrusting the stem of one plant through a hole in a disk of gummed paper so that the paper fits closely against the gills. This disk may be held in place by thrusting the stem of the fungus through a piece of thin paper and then bringing the paper above the cap and twisting the corners. Use white disks for fungi suspected of having coloured spores, and coloured paper for those suspected of

having white spores. If there is any doubt, set two fungi—one with a coloured disk and the other with a white one.

It is an easy matter to fix spores for future use, but a more difficult matter to fix them so that they represent a picture of the radiating gills. In order to do this the stem must be cut from the cap, and the cap must be so arranged over the paper that no draughts shall disturb the spores as they fall, and also so that it may be removed from the paper without disturbing the spores after they have fallen. The writer would suggest that two fine wires should be thrust horizontally and at right angles to each other through the cap, and that the ends of the wires should be supported in a convenient manner, so that the cap may just swing free from the paper which is to receive the spores. The whole must be covered to keep the spores from being disturbed by draughts. When the spores have fallen, the cover can be removed, the cap raised, and the spore-print fixed.

Various methods of fixing spores for prints are recommended. The following are quoted from a bulletin of the Boston Mycological Club :

The following directions for *fixing spore prints* are taken from Herpel ("Das Präpariren der Hutpilze").

Paper which is somewhat absorbent must be used; unglazed blue or black paper (of which the colour must be unaffected by the fixative) for white-spored species. The piece of paper bearing the spore print is to be laid, spores upward, in a flat plate or platter on which a thin layer of fixative has first been poured. The fixative is allowed to soak up through from below, and should not overflow the edges of the paper. When it is certain that the spores as well as the paper are thoroughly soaked, the preparation is removed and dried; sometimes, to prevent sticking, being laid on moistened blotting-paper.

The fixative to be used will vary with the species. For instance, the spores of *Cantharellus cibarius* and some others may be fixed by water alone. The following solution is recommended for *Boleti* and species with coloured spores: One part sandarac, two parts mastic, and two parts Canada balsam, dissolved in thirty parts of ninety-five per cent. alcohol. In the use of this it has been found that the time of soaking necessary to fix the spores is for *Boleti*, two minutes; *Dermini*, *Coprinarii*, *Gomphidius*, *Paxillus*, *Russula*, and *Lactarius*, four to five minutes; pink-spored agarics, also dark brown spored (as the meadow mushroom), and *Cortinarii*, six to eight minutes. It is important that the alcohol should be full strength.

A gelatine solution is useful for white-spored species. This is prepared and used warm. Its strength varies with the species. *Lepiota procera*, *Collybia radicata*, and *Clitocybe laccata* may be fixed by a solution of one part gelatine to thirty of water. For species of *Tricholoma* this is too strong, and one to sixty, or one to two hundred must be the formula. The difficulty arises here from the fact that an excess of

gelatine makes the spores transparent and even invisible. Their opacity may be secured by previous treatment with a solution of one part mastic in thirty of ether.

For certain kinds (*Tricholoma personatum*, *Lepiota granulosa*, *Amanitopsis vaginata*, and others which experiment will discover), ten to twenty-five per cent. of alcohol must be added to the gelatine solution in order to make the spores adhere.

Experience will doubtless show that other fixatives may be used. Gum arabic, for instance, suggests itself. This, however, if strong, is apt to cake the spores together. If one method fails, invention and repeated trials must find a successful means. Reports are requested from all who engage in the amusement of making spore prints. To these a suggestion not without value is that dry agarics (like *Marasmius* and some *Collybias*) may be kept in a condition to shed spores by putting moistened blotting-paper under them.

Another method of making spore prints is to spray them from an atomizer with a solution of white shellac in alcohol. A saturated solution should be made, and then diluted fifty per cent. with alcohol.

THE SEARCH FOR A NAME.—When looking up a name for a plant, the best plan is to use fresh specimens, and, if a good supply can be had, make a careful comparison of all, so as to be sure that the characteristics are normal and not due to injury. If the attempt to find a satisfactory description is not successful, preserve the specimens dried, together with full notes, and send a part of them to the State botanist for determination.

The name of a fungus is not the vital thing. In pursuit of a name do not neglect the plant. Observe it as it grows and in its different stages. Make a friend of it, and you will find it good company.

THE PREPARATION OF ROUGH-DRIED PLANTS FOR THE HERBARIUM—Put the dried plants in a place where they will absorb just moisture enough to make them pliant. Either put them in a box containing something damp, as a wet sponge, sand, or paper; or spread them where they can absorb the moisture of the atmosphere without getting too wet. When pliant, bend the stem and cap so that they lie in the same plane, and arrange them in as natural a form as possible; then place them between driers of unglazed paper, with a weight just sufficient to keep them from curling out of shape.

MOUNTING—The specimens may be placed loose in envelopes made by folding paper as for mosses or lichens, or they may be glued directly to mounting sheets, or they may be kept in boxes of varying sizes.

SECTIONS—A section of a fungus is a very thin slice cut from the plant by running a thin-bladed knife from the top of the cap down through the stem. When well made, sections of young and mature plants are valuable in addition to the notes and dried specimens. A section to be of any value must show the form of the cap; the attachment of the gills to the stem; the thickness of the stem; and the interior, whether solid, hollow, or stuffed. To preserve the section, it must be placed, while fresh, upon a sheet of gummed paper, and then covered with a sheet of waxed paper, and placed between driers, under heavy pressure.

POISONING HERBARIUM SPECIMENS—It will be found necessary to use every means possible to keep insects from the herbarium, as fungi are particularly subject to such pests. In order that no eggs and larvæ may be packed away with the dried plants, it will be well to apply a poisonous solution to the specimens just after they have been moistened to be put into press. Professor Peck, the State botanist of New York, uses a solution made by dissolving strychnine in warm water, and then adding alcohol in sufficient quantities to make the mixture spread easily with a brush.

Sulphate of strychnia	½ ounce
Warm water	4 or 5 ounces
Alcohol	About 2 ounces

In addition to this precaution, the specimens must be kept where insects cannot get at them, or the havoc which they make will be disastrous. An ingenious person can improvise all the apparatus necessary for a successful collection of moderate size; and then, if his enthusiasm continues, he can provide himself with everything of the most improved style from dealers who make a specialty of botanical supplies.

CHAPTER XV : FUNGI FOR THE TABLE

BEFORE you attempt to use fungi for the table be sure that they are edible ; the consequences which follow a mistake are too serious to warrant any risks. Unless you are experienced in making careful observations and comparisons, eat only those fungi which have been shown to you by some one who has tried them and knows them to be wholesome. If you are experienced in making careful observations and comparisons, and wish to make experiments, make them cautiously, using a small quantity of the fungus for the first trial, and, if no ill effects are felt, increase the amount until you are satisfied as to its edibility.

There is no general rule by which one may know an edible species from a poisonous species. One must learn to know each kind by its appearance, and the edibility of each kind by experiment.

Some *edible* mushrooms change colour when bruised, some *edible* ones do not.

Some *poisonous* mushrooms change colour when bruised, and some *poisonous* ones do not.

Some mushrooms with bright colours, or viscid caps, or pleasant taste, or agreeable odour are *edible*, and *some* are *poisonous*.

Some edible mushrooms will turn a silver spoon black, and so will *some poisonous* ones.

Cautions for the Inexperienced

Never use specimens which are decomposed in the *slightest* degree.

Never use those which are at all burrowed by insects.

Never collect, for food, mushrooms in the button stage, as it is difficult for a novice to distinguish the buttons of poisonous species from the buttons of harmless species.

Never use fungi with swollen bases surrounded by sac-like or scaly envelopes.

Never use fungi with milky juice unless the juice is reddish.

Never use fungi with caps thin in proportion to the width of the gills when the gills are nearly all of equal length, especially if the caps are bright coloured.

Never use for food tube-bearing fungi in which the flesh changes colour when cut or broken, nor those with the tubes reddish. Be very cautious with all fleshy tube-bearing fungi.

Never use for food fungi with a web-like ring around the upper part of the stem.

The novice may safely experiment with *Clavarias*, coral fungi, morels, and puffballs.

THE FOOD VALUE OF FUNGI.—Many people thoughtful for the welfare of those with limited opportunities for varying their bill of fare have hoped to solve the problem by introducing into more general use the varied and abundant fungi which grow everywhere throughout our country. In order to accomplish this object, bulletins have been published by the several agricultural departments, and have been distributed freely to those wishing to have them. The result has been that a wide-spread interest has been created in this branch of botany, and fungi have become a common dish on tables where they were never before seen.

As accessories, for relish or variety, edible fungi are undoubtedly valuable; but that they can never take the place of meat, as many fondly hoped, nor rank very high as an essential food, has been shown by the experiments of Mr. L. B. Mendel in the Sheffield Laboratory of Physiological Chemistry, Yale University. Mr. Mendel has demonstrated by chemical analysis and by experiments in artificial digestion that the proportion of proteid matter—the material which meat supplies—is smaller than it was formerly supposed to be, and also that a large proportion of that present is not acted upon by the digestive juices. Since, also, the proportion of water to solid matter is very great, being from seventy to ninety per cent. in the most desirable edible species, it would be necessary that a man should eat a great many pounds of even the richest fungi in order to obtain the daily requisite of proteid matter necessary to maintain a healthy constitution.

The specimens marked edible in this book have been repeat-

edly tried by many people, and without ill effects. Some marked edible are harmless, but poor, while others are extremely delicious and appetizing.

A few directions for preparing different species for cooking are given below, that those who wish to experiment may have the benefit of the experience of others. Receipts for cooking the common mushroom, *Agaricus campestris*, may be found in all complete cook-books, and these receipts as they are given or modified may be used for other kinds also, provided that the directions for the preparation of the different species are followed.

To Keep Mushrooms Temporarily.—Cleanse, remove the parts to be rejected, rinse in cold water the parts to be used, dry with a cloth, then put in boiling water and keep boiling for five or ten minutes. Drain, and wipe dry.

To Prepare the Edible Agarics for Cooking.—Cleanse, cut off the stems and throw them away. Rinse the caps in cold water, drain, and leave in cold water acidulated with lemon or vinegar until just before using.

To Toast Agarics.—Dry with a cloth, dust with flour, put a little butter, pepper, and salt on the gills. Lay the caps, gills upward, on a wire-net toaster, over a moderate fire, and cook from five to ten minutes.

To Bake Agarics.—Dry with a cloth. Line a porcelain pie-dish with toast, spread the peeled caps on the toast, sprinkle with pepper and salt, and pour over them a few spoonfuls of thick cream. Cover with a plate, and place in a moderate oven for fifteen minutes. Serve hot. Or,

Line the dish with toast dipped in hot water and buttered. Spread the caps on the toast, with half a teaspoonful of butter on each one. Cover, and cook in a warm oven for ten minutes.

To Broil Agarics—Broil lightly on both sides over a bright fire. Arrange on buttered toast, sprinkle with pepper and salt. Put bits of butter or bits of toasted bacon on each, and set in the oven a moment. Serve hot.

Mushrooms Stewed—Caps, peeled, one quart; butter, two tablespoonfuls; salt, one teaspoonful; pepper, one-third of a teaspoonful; water or stock, one-half cupful. Boil gently in a covered stewpan for five minutes. Or,

Caps, peeled and cut in pieces, one pint; butter, one table-

spoonful; salt, one-half teaspoonful; pepper, one pinch. Simmer in a stewpan for ten minutes.

To Prepare Russulas—Reject the stems and gills, but not the peel. Cleanse the caps, rinse in cold water, then put for a moment in boiling water, and dry with a cloth. Cook by the receipts given for agarics.

To Prepare Fungi with Milky Juice—Remove the stems, cleanse, rinse, and scald. Steep for six hours in a liquid prepared by mixing one wineglass of strong vinegar, one tablespoonful of salt, and one pint of water. Boil for ten minutes in salt and water. Cook by the receipts given for agarics.

Lactarius deliciosus—Known by its greyish-orange cap marked with brighter zones, and by its orange milk. Also

Lactarius volemus—Known by its reddish-brown cap, two to five inches across, with stems of the same colour, and white milk. May be prepared by simply removing the bases of the stems and then rinsing in water.

To Prepare Amanitas—Reject the stems and peel, and cook but a short time.

To Prepare Chanterelles—Cut off the base of the stems, rinse in cold water, soak in warm milk for six hours. Stew a long time with plenty of butter or stock. Use with meat hashes and stews, or in omelet.

To Fry Chanterelles—Wash, slice, put in melted butter, and stir for ten minutes, simply keeping them warm. Add more butter, pepper, salt, crumbs of bread, and minced parsley, and fry over a hot fire.

To Prepare Coprini—Ink Caps—Use only young specimens. Remove the base of the stems, wipe with a damp cloth. Throw for an instant into boiling water. Fry in boiling butter or lard. Remove from the pan as soon as they break or sink. Serve on toast.

To Prepare Boleti—Remove the tubes with a spoon. Reject the stems.

To Prepare Hydnum repandum—Remove the bases of the stems, and scrape off the spines. They require little cooking.

To Prepare Morels—Cleanse; rinse by shaking them in several waters, or run the cold water from the faucet over them until the pits are thoroughly cleansed. They require to be cooked for a long time.

To Prepare Beefsteak Fungus—Gather when of a light-red colour. Remove the hard base, cleanse in cold water.

For salad—Cut in thin slices and serve with dressed lettuce.

Minced—Mince fine, put in a stewpan with butter, three ounces to the pound. Season with salt, pepper, minced parsley, and onion juice. Stew gently for twenty minutes. Bind with egg-yolk beaten in cream, and serve with toast. This mince may be used with veal or chicken hash.

To Prepare Gyromitras.—Cleanse, cut in slices, boil in water fifteen minutes, then wash by shaking in two successive waters boiling hot. Dry on cloths, and cook as directed for morels.

To Prepare Woody Pore-bearing Fungi—Polyporæ.—Take the soft parts of young specimens. Put in boiling water for a few moments, rinse in cold water, and dry on cloths. Spread with butter, lay in a stewpan, and cover; then keep them for ten minutes just warm enough to melt the butter. Strain, broil for fifteen minutes, or stew half an hour or more with gravy.

To Prepare Clavarias and Branched Hydnums.—Cleanse, throw into scalding water for a moment, and then put into cold water made acid with lemon or vinegar until they are to be cooked. Divide the large ones, and tie the small ones into bundles. Place in a stewpan with bits of butter laid on them. Cover the pan, and expose to heat enough to melt the butter. Leave for ten minutes, and drain.

To Cook Clavarias.—Put into a hot stewpan with bits of butter; season with salt, pepper, and lemon juice. Cover closely, and stew for half an hour. Thicken with cream and flour, season to taste, and cook until tender.

Clavarias may also be cooked as directed for agarics.

To Prepare Puffballs.—Cleanse, peel, trim off the base. Cut in two pieces, and reject all those which are not pure white within.

To Cook Puffballs.—Fry in lard five or six minutes, with bacon, parsley, onion juice, salt, and pepper; or cook as directed for agarics.

To Cook the Giant Puffball.—Cut in slices half an inch thick, dip in the beaten yolk of egg, pepper, and salt. Fry in boiling fat for five or six minutes.

Puffball Salad.—Cut in strips, and serve with green salad dressed with mustard, oil, and vinegar.

AUTHORITIES CONSULTED

ATKINSON. Studies and Illustrations of Mushrooms, I, II. Bull. Cornell Univ. Expt. Station, **138**: 337–366, *f. 87–112.* 1897. **168**: 491–516, *f. 83–97.* 1899.

BURNAP. Notes on the Genus Calostoma. Bot. Gaz., **23**: 180–192, *Pl. 19.* 1897.

BURT. A List of the Vermont Helvellaceæ, with Descriptive Notes. Rhodora, 1: 59–67, *Pl. 4.* 1899.

—— On Collecting and Preparing Fleshy Fungi for the Herbarium. Bot. Gaz., **25**: 172–186, *Pl. 14.* 1898.

—— The Phalloideæ of the United States, I–III. Bot. Gaz., **22**: 273–292, *Pl. 11, 12;* 379–391. 1896. **24**: 73–92. 1897.

COOKE. Edible and Poisonous Mushrooms. 8vo, *18 plates.* London, 1894.

—— Illustrations of British Fungi. 8 vols., *Pl. 1–1198.* London, 1881–1891.

—— Mycographia. *Pl. 1–113.* London, 1879. [Pezizales.]

—— Mycographia. 1–10, 87–104, 179–206, 215–220; *Pl. 1–4, 41–46, 81–96, 101, 102.* 1879. [Helvellales.]

COVILLE. Observations on Recent Cases of Mushroom Poisoning in the District of Columbia. U. S. Dep. Agr., 1897.

FALCONER. Mushrooms: How to Grow Them. U. S. Dep. Agr., 1896.

FARLOW. Edible and Poisonous Mushrooms. U. S. Dep. Agr., 1898.

FRIES. Icones selectæ Hymenomycetum, Holmiæ. 2 vols. 1867–1884.

HAY. Text-book of British Fungi. 8vo. London, 1887.

MASSEE. A Monograph of the Geoglosseæ. Annals of Bot., 225–306, *Pl. 12, 13.* 1897.

—— A Monograph of the Genus Calostoma. Bot. Gaz., **23**: 180–192, *Pl. 19.* 1897.

—— A Monograph of the British Gastromycetes. Annals of Bot., Vol. IV, No. XIII. November, 1889.

MENDEL. The Chemical Composition and Nutritive Value of some Edible American Fungi. Am. Jour. of Physiology, **1**: 225–238. 1898.

MICHAEL. Führer für Pilzfreunde. 12mo, *55 plates*. Zwickau, 1897.

MORGAN. North American Fungi. Jour. Cincinnati Soc. Nat. Hist., II: 141–149, *Pl. 3*. 1889. [Phallales.]

—— North American Fungi. Jour. Cincinnati Soc. Nat. Hist., **12**: 8–22, *Pl. 1, 2*. 1889. 163–172, *Pl. 16*. 1890. **13**: 5–21, *Pl. 1, 2*. 1891. **14**: 141–148. *Pl. 5*. 1892. [Lycoperdales.]

PECK. Reports of the State Botanist of New York, in the Regents' Reports of the State Museum of Natural History, 22–51.

—— Boleti of the United States. Bull. N. Y. State Mus., No. 8. 1888.

STEVENSON. British Fungi. 2 vols.

UNDERWOOD. Moulds, Mildews, and Mushrooms. New York, 1899.

—— On the Distribution of the North American Helvellales. Minn. Bot. Studies, I: 483–500. 1896.

WEBSTER. Notes on Calostoma, Rhodora, I: 30–33. 1899.

NOTE

It is customary, when writing the name of a fungus for scientific purposes, to append the name of the author who first published the appellation. The author's name, for convenience, may be abbreviated. A list of such abbreviations as are used in this book is given below.

A. & S. Albertini and Schweinitz.

BATSCH. Augustus Batsch (1761–1802), German botanist.

BERK. Rev. Miles Joseph Berkeley.

B. & C. Berkeley and Curtis.

BOSC. Louis Bosc (1759–1828), one of the first collectors of fungi in the United States.

BULL. Pierre Bulliard, 1742–1793.

BURNAP. Charles E. Burnap, an American student.

BUXB. Johann Christian Buxbaum, 1693–1730.

D. C. Augustin Pyrame de Candolle (1778–1841), a prominent Swiss botanist.

DESV. Nicaise Augustin Desvaux, French botanist, 1784–1856.

ELLIS. J. B. Ellis. Mr. Ellis is a mycologist in the United States. The Ellis collection of fungi contains the largest number of types of any collection of American fungi in existence. It is deposited in the museum of the New York Botanical Garden.

FR. Elias Magnus Fries (1794–1878), a Swedish botanist, who laid the foundations for the study of the Basidiomycetes.

HOLMSK. Theodor Holmskiold (1732–1794), a Danish botanist.

HUDS. William Hudson (1730–1793), an English botanist.

LASCH. Wilhelm Lasch (1786–1863), a German botanist.

LK. Heinrich Friedrich Link (1767–1851), a German botanist.

L. or LINN. Carl von Linnæus (1707–1778), a Swedish botanist, who revised the principles of classification and introduced what is known as the *binomial nomenclature*. According to his method, the name of a plant is reduced to two words: the first, or

generic, name is a substantive or a word used as a substantive ; while the second, or specific, name is an adjective. *Lactarius* is the generic name of those fungi dripping milk, and *deliciosus* (delicious) the specific name for one edible species.

MASS. George Massee, an English botanist.

MORG. A. P. Morgan, an American botanist.

PK. Charles H. Peck (1833–), the State botanist of New York.

PERS. Christian Hendrik Persoon (1755–1837), a German botanist.

ROZE. Ernest Roze, a French botanist.

SCHW. Lewis David de Schweinitz (1780–1834), an American botanist ; one of the first to make mycology a serious study.

SCOP. Giovanni Antonio Scopoli (1723–1788), an Italian botanist.

SCHÆFF. Jacobi Christiani Schæffer (1718–1790), a German botanist.

SACC. P. A. Saccardo (1845–), an Italian botanist. Saccardo is the compiler of *Sylloge Fungorum*, a work in Latin, containing descriptions of over forty thousand species. It is a most valuable work, as it has made accessible to workers throughout the world the greater part of the technical descriptive literature upon the subject of fungi.

VAHL. Martin Vahl (1749–1804), a Norwegian botanist.

The diacritical marks used in the pronunciation of the Latin names indicate the sounds of the same letters in the following list :

ă	as in	fat	ŭ	as in	cut
ā	"	fate	ū	"	cūte
ĕ	"	met	o͝o	"	mo͝oth
ē	"	meet	o͞o	"	mo͞o
ĭ	"	fin	ġ	"	gem
ī	"	fine	g	"	get
ŏ	"	not	c	"	cat
ō	"	note	ç	"	cent

INDEX AND GLOSSARY

Adnate, 35; united by growth; said of gills when they are grown to the stem.
Agaricaceæ, 30, 32, 40, 46, 73.
Agaricales, 14, 30.
Agaricus, 39, 73.
— abruptus, 77.
— arvensis, 75.
— campestris, 12, 75, 76.
— cretaceus, 64.
— hemorrhoidarius, 75.
— how to grow, 73.
— maritimus, 75.
— placomyces, 75.
— Rodmani, 75, 76.
— subrufescens, 75.
— sylvaticus, 75.
Algæ, 9.
Algal-like fungi, 9, 17.
Amanita, 34, 46.
— Cæsarea, 50, 52.
— Frostiana, 52.
— muscaria, 52.
— phalloides, 48.
— to prepare, for cooking, 154.
— verna, 65.
— young plant, 47.
Amanitopsis, 34, 53, 88.
— parcivolvata, 55.
— vaginata, 54.
— — var. alva, 54.
— — var. fulva, 54.
— — var. livida, 54.
Anthurus, 26.
— borealis, 121.
Antidote, 48; anything that will counteract the effects of poison.
Arachnoid, 38; web-like.
Armillaria, 34.
— mellea, 61.
— — var. bulbosa, 62.
— — var. exannulata, 62.
— — var. flava, 62.
— — var. glabra, 62.
— — var. obscura, 62.
— — var. radicata, 62.
Asci, 18, 19; the sacs in which the spores of the cup fungi are developed.
Ascoma, 19, 20, 21; the disk-like body bearing the spore-sacs of the cup fungi and their allies.
Ascomata, 20, 21; plural of ascoma.
Ascomycetes, 17, 18, 135.
Ascus, 17; singular of asci.
Astræus hygrometricus, 130.

Basidia, 16, 17; enlarged cells bearing spores.
Basidiomycetes, 14, 17, 97, 116.
Bean seed to show embryo, 7.
Bear's head, 2, 96.
Beefsteak fungus, 103.
— for salad, 155.
— minced, 155.
— to prepare, 155.
Bird's-nest, 24, 133.
Bitter Boletus, 107.
Black knot, 9.
Boletaceæ, 31, 44, 102.
Boleti, 14.
Boleti, care of spores, 148.
— to prepare, 154.
— of the United States, 104.
— which change colour, 104.
Boletinus, 45, 102, 103.
— pictus, 103.
— painted, 103.
Boletus, 44, 104.
— Americanus, 109.
— bicolor, 105.
— calopus, 106.
— castaneus, 108.
— chrysenteron, 106.
— cyanescens, 105.
— edulis, 109.
— eximius, 108.
— felleus, 107.
— glabellus, 104.
— mutabilis, 105.
— pallidus, 105.
— Peckii, 106.
— purpureus, 107.
— radicans, 106.
— Satanus, 107.
— scaber, 108.
— speciosus, 105.
— subtomentosus, 109.
— versipellis, 108.
Boston Mycological Club, 145.
Bovista, 128.
— plumbea, 128.
— section of, 128.
Bovistella, 29, 124, 129.
— Ohiensis, 129.
Brackets, 2, 4.
Brain-shaped Calvatia, 126.
Branches, 7.

Brick top. (Pl. opposite p. 81.)
Brick-red mushroom, 80.
Button, 11, 12; the name given to a young mushroom.

Cæsarea, 50.
Cæsar's mushroom, 51.
Cæspitose, 145; growing in clumps, many plants from one root.
Calostoma, 29, 123, 124, 131.
— cinnabarinum, 132.
— lutescens, 132.
— Ravenelii, 132.
Calvatia, 28, 124, 126.
— craniformis, 126.
— cyathiformis, 128.
— fragile coat, 126.
— maxima, 127.
— section of, 126.
Cambium, 137; the delicate young cells between the wood and bark. The inner become new wood; the outer, new bark.
Campestris, 76.
Caninus, 120.
Cantharellus, 36, 53.
— cibarius, 148.
— floccosus, 53.
Cap, defined, 12.
— membranous, 35.
— ragged edge, 13.
— scaly, 34.
— smooth, 34.
— striate, 50.
— umbonate, 63.
Capillitium, 25, 126: simple or branched threads mixed with spores.
— elastic, 126.
Capitate, 21; having a distinct head.
Capitila. (Pl. opposite p. 136.)
Caput-ursi, 96.
Carbon, 4; an elementary substance found in plant and animal tissue.
Cartilaginous, 36, 146; similar to the elastic tissue found at the ends of fresh bones.
Caterpillar, fungus growing on, 136.
Cautions, 151.
Cell, 11; a living vegetable unit.
— spore, 11.
— club-like, 13, 14.
Cervinus, 87.
Chalk agaric, 64.
Chanterelle hygrophorus, 59.
— to fry, 154.
— to prepare, 154.
Chemical analysis, 152.
Chestnut Boletus, 108.
Chitonia, 83.
Chlorophyll, 6, 144; the green coloring matter of plants.
Christmas greens, 5, 7.
Chrysenteron, 106.
Cibus Deorum, 51.
Cinnabarinum, 132.
Circinatus, 113.
Class, 7, 15, 17.
Classification, 5, 7.
Clathraceæ, 26, 117, 121.
Clathrus, 26, 121.
— cancellatus, 121.
— columnatus, 121.
Clavaria, 3, 14, 99.
— aurea, 100.
— botrytes, 100.
— cristata, 100.
— fellea, 101.
— flava, 99.
— formosa, 101.
Clavaria, golden, 100.
— pale yellow, 99.
— pistil, 101.
— pistillaris, 101.
— red-tipped, 100.
— to cook, 155.
— to prepare, 155.
Clavariaceæ, 30, 98, 99.
Clavate, 21; club-shaped.
Clitocybe, 35, 68, 70.
— laccata, 70.
— care of spores, 148.
Clitocybe virens, 71.
— var. amythystina, 71.
— var. pallidifolia, 71.
— var. striatula, 71.
Clitopilus, 37, 88.
Collector, notes of, 145.
— outfit of, 147.
Collybia, 35, 66.
— care of specimens, 150.
— care of spores, 148.
— familia, 67.
— radicata, 66.
— radicata, care of spores, 148.
— velutipes, 66.
Columella, 125.
Common mushroom, 76.
Context, 42; the tissue immediately in contact with a specified part.
Coprinarii. See Coprini.
Coprini, care of spores, 148,
— to prepare, 154.
Coprinus, 40, 89.
— atramentarius, 91.
— comatus, iii, v, viii, 90.
— micaceus, 91.
Coral fungi, 98.
Coral Hydnum, 97.
Cordyceps, 136.
Corollas, 6.
Cortinarii, care of spores, 148.
Cortinarius, 38, 85.
— alboviolaceus, 87.

Cortinarius violaceus, 87.
Count de Vecchi, 49.
Craniformis, 126.
Craterellus cantharellus, 73.
Crested Clavaria, 100.
Crucibulum, 133.
Cup fungi, 19, 137.
Cup-shaped puffball, 128.
Cyathiformis, 128.
Cyathus, 133.
Czar Alexis, 49.

Dædalea, 43.
Death Cup, 48.
Deconica, 39, 82.
Decurrent, 35; growing down the stem; said of gills.
Deliquescent, 26; dissolving or melting away.
Destroying Angels, 2. See Death Cup.
Dictyophorus, 27.
Dissemination, of seeds, 6; the act of scattering.
— of spores, 4, 118, 126, 131, 133, 137.

Early Pholiota, 84.
Earth-stars, 123, 129.
Earth-tongues, 138.
Eccilia, 87, 88.
Edible Boletus, 109.
Elaphomyces, 136.
Elias Fries, 97.
Elm Pleurotus, 58.
Emetic Russula, 68.
Entoloma, 37, 88.

Fairy clubs, 9.
Fairy-ring mushroom, 66.
Fairy rings, 3.
Families, 7, 16.
Fat Pholiota, 83.
Favolus, 115.
— areolarius, 115.
Fawn-coloured Pluteus, 87.
Fern with spore capsules, 8.
Fertile gleba, 125.
Fibrillose, 146; furnished with fibres.
Fistulina, 44, 102.
— cavipes, 103.
— decipiens, 103.
— hepatica, 102.
— paluster, 103.
— porosus, 103.
Fistulose, 146; hollow through the whole length.
Flexuous, 146; bent alternately in opposite directions.
Flint-stone, 127; a stone formerly used for obtaining sparks.
Floccose, 45; clothed with locks of soft hair or wool.
Flocculent, 52; woolly.
Flowerless plants, 5.
Fly Amanita, 49.
Food of the gods, 51.
Frog-spittle, 6.
Fruiting portion, 12, 13.
Führer für Pilzfreunde, 104.
Fungi, defined, 4.
— for food, 1, 151.
— for the herbarium, 145.
— for the table, 151.
— not grewsome, 2.
— habitat, 2.
— how to grow them, 73.
— number of, 15.
— poisonous, 1, 48, 151.
— typical parts, 12.
— with gills, 30, 32–40, 46–92.
— with milky juice, 92, 93.
— with pores, 42, 44, 102, 155.
— with spines, 23, 41, 94–97.
Fungi with teeth, 94–97.
Fungus plant, 9.
Fusiform, 40; cylindrical, tapering gradually to each end.

Gasteromycetes, 2, 3, 14, 24.
Geaster, 28, 124, 129.
— hygrometricus, 130.
— minimus, 130.
Gelatinous fungi, 116.
Genera, 7, 16.
Geoglossaceæ, 21, 138.
Geoglossum, 139.
— glabrum, 139.
— hirsutum, 139.
Giant puffball, 127.
Gills, 12.
— forked, 53.
— notched, 58.
— structure, 13.
— toothed, 56.
— waxy, 61.
Glabrous, 50; not hairy.
Gleba, 26, 117; the spore mass of the pouch fungi.
Glistening Coprinus, 91.
Globose, 40; having or approaching a spherical form.
Golden Clavaria, 100.
Golden-flesh Boletus, 106.
Golden Peziza, 138.
Gomphidius, 4, 92.
— fixing spores of, 148.
Grandinia, 94.
Grass spikelet, 7.
Gray-gilled mushroom, 79.
Green Russula, 69.
Greeks, 1.
Guepinia, 116.
Gyromitra, 20, 140, 141.
— esculenta, 141.
— to prepare, 155.

Habitat, 145; the locality where a plant naturally lives.
Haschisch, 49; an intoxicating preparation of the Indian hemp.
Hedgehog Hydnum, 97.
— mushroom, 94.
Helvella, 20, 140.
— elastica, 142.
— lacunosa, 143.
— — diagrammatic drawing, 143.
— — section of stem, 143.
Helvellaceæ, 20, 140.
Helvellales, 19, 138.
Hirneola auricula-Judæ, 116.
Honey-coloured Armillaria, 61.
Host, 22; the plant or animal which supports a parasite.
Hydnaceæ, 32, 94.
Hydnum, 14, 32, 41, 94, 116.
— albidum, 95.
— branched, to prepare, 155.
— caput-medusæ, 97.
— caput-ursi, 96.
— coralloides, 97.
— echinaceus, 97.
— imbricatum, 96.
— repandum, 95.
— rufescens, 95.
Hydrogen gas, 4; one of the elements found in plant and animal tissue.
Hygrophanous, 145; opaque when dry, and transparent when moist.
Hygrophorus, 35, 59.
— cantharellus, 59.
— eburneus, 61.
— miniatus, 60.
Hygrophorus, var. flava, 60.
— var. flaviceps, 60.
— var. flavipes, 60.
— var. rosea, 60.
Hymenogastrales, 133.
Hymenomycetes, 2, 14.
Hypha, 11; a thread-like strand of the vegetative part of a fungus.
Hypholoma, 39, 78.
— capnoides, 79.
— elæodes, 80.
— epixanthus, 79.
— fasciculare, 80.
— incertum, 80, 81.
— perplexum, 78, 79.
— sublateritium, 81.
Hypocreales, 18, 136.

Imperial mushroom, 51.
Incertum, 80, 81.
Indian pipe, 7.
Ingenious stamens, 6.
Ink-caps, 89.
Inky Coprinus, 91.
Irpex, 94.
Ivory Hygrophorus, 61.

Jew's Ear, 116.
Judas's Ear, 116.
Juice, milky, 32.
— watery, 32.
Juvenal, 1; a Roman poet, first century A.D.

Kaiserling, 51.
Key, 15–45.

Lachnocladium, 99.
Lactarius, 32, 92.
— care of spores, 148.
— channeled, 33.
— crisped, 33.
— deliciosus, 154.
— entire, 1.
— ligniotus, 93.
— piperatus, 92.
Lactarius, to cook, 154.
— to prepare, 152.
— volemus, 154.
Lamellæ, 12.
— acute 35; the edges thin, not blunt.
— adnate, 35; grown to the stem.
— brittle, 35.
— decurrent, 35; growing down the stem.
— entire, 47.
— free from the stem, 34.
— obtuse, 36.
— shallow folds, 36.
— simple, 33.
— sinuate, 35.
— splitting deeply, 33.
— toothed, 32.
— villous, 33; with soft hairs.
— waxy, 35.
Larch canker, 137.
Large club, 101.
Latticed Clathrus, 121.
Leaf-green, 4, 6, 15, 144.
Lentinus, 32.
— lepideus, 56.
Lenzites, 43.
— betulina, 113.
— separia, 114.
Lepiota, 34, 63.
— Friesii, 65.
— granulosa, care of spores, 149.
— naucinoides, 64.
— procera, 63.
— — care of spores, 148.
Leptonia, 88.
Light from dry rot, 2.
Liverworts, 5, 7.
Lycogala epidendron, 144.
Lycoperdales, 25, 28, 123.
Lycoperdon, 29, 124.
— cyathiforme, 128.
— giganteum, 127.
— maxima, 127.
— pyriforme, 125.

Lycoperdon, section of, 124.
— subincarnatum, 125.
Lycopodiums, 7.

Majoon, 49; an intoxicating confection of India.
Maned agaric (see Comatus), ii, v, viii, 90.
Marasmius, 33, 65.
— care of spores, 149.
— oreades, 66.
Marchantia polymorpha, 7.
Masked Tricholoma, 72.
Medusa's head, 97.
Mendel, 152.
Membrane fungi, 14, 22.
Merulius, 110.
— lacrymans, 110.
Mitrula, 21.
— vitellina, var. irregularis, 140.
Morels, 140, 154.
— to prepare, 154.
Morchella, 21, 140, 141, 142.
— angusticeps, 142.
— bispora, 142.
— deliciosa, 142.
— esculenta, 141, 142.
— section of, 141.
— semilibera, 142.
Mould, 2.
— on bread, 9.
— on food, 2.
— on dead fly, 9.
Mould-like fungi, 15, 77.
Mounting, 149.
Mount Marcy, 68.
Mucronella, 94.
Muscaria, 49.
Muscarine, 50; a poison found in A. muscaria.
Mushrooms, baked, 153.
— broiled, 153.
— stewed, 153.
— toasted, 153.
Mushrooms, to keep, 15.
— to prepare, 153.
Mutinus, 27.
— bambusinus, 120.
— caninus, 120.
— embryo plant, 120.
— young plant, 120.
Mycelial threads, 117, 123.
— luminosity of, 63.
Mycelium, 11, 110, 136.
— food provider, 13.
Mycena, 35, 55.
— hæmatopoda, 55.
Myxomycetes, 144.

Name, not a vital thing, 150.
— the search for, 150.
Nidularia, 133.
Nidulariales, 24, 133.
Nolanea, 88.

Odontium, 94.
Offensive fungi, 117.
Omphalia, 35, 68.
— umbellifera, 68.
Orange Amanita, 50.
Orange-cap Boletus, 108.
Orders, 7, 16.
Oreades, 3.
Oxygen, 4; one of the elementary substances which, chemically united with carbon and hydrogen, forms plant tissues.
Oyster mushroom, 57.

Painted Boletinus, 103.
Panæolus, 92.
Panus, 33, 67.
Paraphyses, 135; jointed thread-like bodies found accompanying the spore-sacs of certain fungi.
Parasite, 22; that which lives by taking its food from living plants or animals.
Parasite, Cordyceps, on Elaphomyces, 136.
Parasol mushroom, 63.
Paxillus, care of spores, 148.
Peach cure, 9.
Pear-shaped puffball, 125.
Peridium, 14, 23, 24, 25, 123; the thickened covering to a puffball.
Perithecium, 58; a rounded, oval, pear-shaped, or beaked body in which the spore-sacs are developed.
Perplexing Hypholoma, 78, 79.
Perplexum, 78.
Peziza, 22, 137.
— æruginosa, 137.
— aurantia, 138.
— odorata, 137.
— Willkommii, 137.
Pezizales, 19, 20, 137.
Phallaceæ, 27, 117.
Phallales, 24, 117, 121.
Phallin, 48; a deadly poison found in certain fungi.
Phalloides, Amanita, 48.
Phallus, 27.
— impudicus, 117.
— section of young impudicus, 119.
Phlebia, 94.
Pholiota, 38.
— adiposa, 83.
— limonella, 83.
— præcox, 84.
Phycomycetes, 9, 17.
Phylæ, 7.
Physalacria, 98.
Pileus, 146; the cap of a mushroom.
Pilosace, 39, 82.

Pinkish puffball, 125.
Pistachio nuts, 78.
Pistil of St. Johnswort, 6.
— of violet, 6.
Pistil Clavaria, 101.
Pistillaria, 98.
Pleurotus, 32.
— ostreatus, 57.
— sapidus, 58.
— ulmarius, 58.
Pliny, 135.
Pluteus, 37.
— cervinus, 87.
Poison Amanita, 48.
Poisoning, cases of, 49.
— of herbarium sheets, 150.
— recipes for, 150.
Poisonous, 48, 49, 151.
Pollen, 4, 6.
Polypodium vulgare, 6.
Polyporaceæ, 31, 42, 102, 109.
Polypori, 110.
— to prepare, 155.
Polyporus, 42.
— arcularius, 112.
— circinatus, 113.
— conchatus, 111.
— fomentarius, 110.
— megaloma, 110.
— perennis, 111.
— pergamenus, 111.
— squamosus, 112.
— sulphureus, 111.
— velutinus, 111.
Pores, 14; the openings of the spore-bearing tubes.
— easily separating from the adjoining tissue, 44, 102.
— in radiating rows, 102.
— in the form of tubes whose mouths are separated, 44, 102.
— long-hexagonal, 43.
Pores, permanently united to adjoining tissue, 42, 102.
— separating with difficulty, 45.
— to prepare woody, 155.
Pouch fungi, 14, 23, 123.
Pruinose, 146; as if frosted with a bloom or powder.
Psathyra, 39, 82.
Psathyrella, 40, 92.
Psilocybe, 39, 82.
Protoplasm, 144; the viscid, contractile, semi-fluid substance of an animal or vegetable cell.
Protozoa, 144; unicellular microscopic animals.
Pterula, 99.
Puffballs, 3, 4, 123.
— salad, 155.
— to cook, 155.
Purple Boletus, 107.
Pycnodon, 94.

Radulum, 94.
Resupinate, 31.
Ring, 12.
— movable, 63.
Rodman's mushroom, 76.
Romans, 1.
Rough-dried plants, 149.
Russia, 49.
Russula, 3, 35.
— care of spores, 148.
— emetica, 68.
— heterophylla, 70.
— to prepare for cooking, 154.
— virescens, 69.

Salmon fungus, 9.
Saprophytes, 123.
Scaly Lentinus, 56.
Schizophyllum, 33, 67.
Scleroderma, 133.
Scleroderma vulgare, 134.
Sclerodermatales, 25, 133.
Seed-boxes, 6, 8.
Sensitive plant, 89.
Shaggy-mane, ii, v, viii, 90.
Shelley, 1, 2, 89.
Silver fir, 6.
Simblum, 26.
— rubescens, 122.
Sinuate, 35; said of gills which have a notch or recess near the stem.
Slime fungi, 144.
Smallest Earth-star, 130.
Smith, Rev. Gerard, 3.
Smokeballs, 123.
Smooth Lepiota, 64.
Smuts, 2, 14, 22.
Sparassis crispa, 99.
Spathularia, 21, 138.
— velutipes, 138.
Species, 7, 16.
Specimens, 7.
— care of, 147.
Spencer, 2, 5.
Sphæriales, 136.
Sphærobolus, 133.
Spines, 14, 41.
Spore-dust to stanch blood, 127.
Spore print, 4.
— receptacle, 10.
Spore-sac fungi, 9, 17, 18, 135.
Spores, 11; single cells which serve the same purpose for fungi as seeds do for flowering plants.
— black, 32, 46, 89–92.
— brown, 38, 73–83.
— colour, 4.
— collecting of, 147.
— dark brown, 32, 73–83.
— dissemination of, 4, 118, 126, 131, 133, 137.
— elliptical, 72.
— fusiform, 40.

Spores, how formed, 12.
— in a definite rind, 14, 23, 123.
— on spicules, 9.
— pink, 36, 46, 87, 88.
— purplish-brown, 32.
— rough, 69.
— rosy, 32, 87, 88.
— rusty brown, 32, 46, 83–87.
— salmon, 32, 87, 88.
— violet, 32, 58.
— white, 32, 46–73.
— yellowish-brown, 32, 83–87.
Spreading Hydnum, 195.
State botanist, 150.
Stem, 12, 13.
— cartilaginous, 35.
— central, 60.
— characters of, 46.
— cup at base, 47, 49, 50, 54, 88.
— eccentric, 56.
— fleshy, 5.
— hollow, 50.
— lateral, 58.
— mealy, 46.
— rusty, 46.
— smooth, 46.
— solid, 56.
— swollen base, 48.
— wanting, 113.
Sterigmata, 9, 17; the slender stalks or spicules upon which the spores of the Basidiomycetes are borne.
Sterile, 29; not containing spores.
— base, 125.
Stinkhorns, 24, 26, 117, 123.
Stipe, 12; stem of fungus.
Striate, 35, 50; said of the cap when marked with radiating straight lines on the margin.
Strobilomyces, 102.
Stroma, 136; a compact mass of mycelium which unites the pear-shaped bodies in which spore-sacs are contained.
Stropharia, 39, 82.
Sulphury polyporus, 111.
Sistotrema, 94.

Tall Lepiota, 63.
Thelephoraceæ, 73.
Tinder, 127; dry material which will ignite without explosion when in contact with a spark.
Toadstool, 2, 8.
Tox-albumins, 48; poisonous proteids produced by bacteria.
Trama, 42; a middle tissue between two layers of spore-bearing tissue.
Trametes, 42, 110, 113.
— cinnabarina, 113.
— pini, 113.
— suaveolens, 113.
Tremellines, 22.
Tremellodon, 116.
Tricholoma, 35.
— care of spores, 148.
— personatum, care of spores, 149.
Tricholoma personatum, var. bulbosum, 72.
Trogia, 33.
Truffles, 18.
Tuberales, 18, 30, 135.
Tufted yellow mushroom, 80.
Tunbridge ware, 137.
Typhula, 98.

Umbilicate, 146; provided with a central depression.
Umbonate, 63; provided with a central prominence.
Uncertain Hypholoma, 8.

Variable Russula, 70.
Veil, 12, 113; the membrane which covers the spore surface of a fungus in the young stage.
Velvety Spathularia, 138.
Verdette, 69.
Vermilion Hygrophorus, 60.
Vernal Amanita, 65.
Vibrissea, 139.
— circinans, 140.
— truncorum, 139.
Volva, 34, 46, 52.
— of mutinus, 120.
Volvaria, 36, 88.

Water-measuring Earthstar, 130.
White Hydnum, 95.
Wrapper, 47, 49, 50, 51, 54.

Xylaria, 136.

Yeast plant, 9.

INDEX OF PLATES

Agaricus abruptus, 74, 77.
— campestris, 74, 75.
— Rodmani, 76.
Amanita Cæsarea, 50.
— muscaria, 49.
— phalloides, 48.
Amanitopsis parcivolvata, Frontispiece.
— strangulata, 53.
— vaginata, 54.
Armillaria mellea, 61.

Bear's head, 96.
Bird's-nest, 130.
Bitter Boletus, 107.
Boletinus pictus, 103.
Boletus chrysenteron, 60.
— felleus, var. obesus, 107.
— scaber, var. niveus, 108.
Bovistella Ohiensis, 128.
Brain puffball, 126.
Brick top Hypholoma, 81.
Bristly panus, 145.
Broad-gilled Collybia, 66.

Calostoma cinnabarina, 132.
— lutescens, 132.
— Ravenelii, 132.
Calvatia craniformis, 126.
— cyathiformis, 128.
Cantharellus floccosus, 136.
Chanterelle Hygrophorus, 60.
Clavaria aurea, 100.
— formosa, 101.
— ligula, 98.
Clitocybe illudens, 70.
— laccata, 67.
— virens, 71.
Collybia familia, 67.
— maculata, 66.
— platyphylla, 66.
Cone-like Boletus, 102.
Coprinus atramentarius, 91.
—— var. silvestris, 89.
Coral Hydnum, 97.
Cordyceps capitala (Parasitic on Elaphomyces), 136.
Cortinarius alboviolaceus, 65.
— armillatus, 86.
— caninus, 85.
Craterellus cantharellus, 73.
Cup-shaped Calvatia, 128.
Cyathus vernicosus, 130.

Dædalea quercina, 114.
Death cup, 48.
Deceiving clitocybe, 70.
Delicious morel, 142.
Dog cortinarius, 85.

Early Pholiota, 84.
Elaphomyces, 136.
Elfingia fomentaria, 110.
Emetic Russula, 68.

Fat Pholiota, 61.
Fawn-coloured Pluteus, 87.
Field mushroom, 75.
Flesh-coloured puffball, 134.
Floccose Chanterelle, 136.
Fly Amanita, 49.
Fomes fomentarius, 110.

Geaster hygrometricus, 130.
— minimus, 130.
Golden Clavaria, 100.
— flesh Boletus, 60.
— Peziza, 138.
Grainy Lepiota, 63.
Green Russula, 69

Hard-skinned puffball, 134.
Helmet Mycena, 55.
Helvella elastica, 140.
— lacunosa, 140.
Hirneola auricula-Judæ, 116.
Honey-coloured Armillaria, 61.
Hydnum caput-ursi, 96.
— coralloides, 97.
— repandum, 103.
Hygrophorus cantharellus, 60.
— eburneus, 84.
— miniatus, 60.
Hypholoma incertum, 80.
— perplexum, 78.
— sublateritium, 81

Index of Plates

Inky Coprinus, 91.
— — wood variety, 89.
Ivory Hygrophorus, 84.

Jelly-like Tremellodon, 137.
Jew's ear, 116.

Lactarius ligniotus, 93.
— piperatus, 92.
Least Earth-star, 130.
Lentinus lepideus, 56.
— strigosus, 145.
Lenzites betulina, 113.
Leotia lubrica, 137.
Lepiota Friesii, 65.
— granosa, 63.
— nancinoides, 64.
— procera, 64.
Little-tongue Clavaria, 98.
Lycogola epidendron, 136.
Lycoperdon pyriforme, 125, 134.
— subincarnatum, 134.

Masked Tricholoma, 72.
Mitrula vitellina, var. irregularis, 140.
Morchella deliciosa, 142.
Mutinus caninus, 136.
Mycena galericulata, 55.
— hæmatopoda, 93.

Orange amanita, 50.
Oyster mushroom, 58.

Painted Boletinus, 103.
Panus strigosus, 145.
Parasol mushroom, 64.
Pear-shaped puffball, 125.
Peppery Lactarius, 92.
Perplexing Hypholoma, 78.
Peziza aurantia, 138.
— odorata, 138.
Phallus impudicus, 119.
Pholiota adiposa, 61.
— aggericula, 73.
— præcox, 84.
Pleurotus ostreatus, 58.
Pluteus cervinus, 87.
Poison Amanita, 48.
Polyporus arcularius, 142.
— circinatus, 112.
— fomentarius, 110.
— versicolor, 112.

Rodman's mushroom, 76.
Russula emetica, 68.
— virescens, 69.

Scabrous-stemmed Boletus, 108.
Scaly Lentinus, 56.
Scleroderma vulgare, 134.
Sheathed Amanitopsis, 54.
Slippery Leotia, 137.
Smooth Lepiota, 64.
Spathularia velutipes, 132.
Spotted Collybia, 66.
Spreading Hydnum, 103.
Strangled Amanitopsis, 53.
Strobilomyces strobilaceus, 102.

Tall Lepiota, 64.
Tremellodon gelatinosum, 137.
Tricholoma personatum, var. bulbosum, 72.

Uncertain Hypholoma, 80.

Vermilion Hygrophorus, 60.

Water-measuring Earth-star, 130.

Xylaria, 116.

Young pear-shaped puffball, 134.

Zoned Cortinarius, 86.



A FINE IS INCURRED IF THIS BOOK IS NOT RETURNED TO THE LIBRARY ON OR BEFORE THE LAST DATE STAMPED BELOW.

JUN 7 1973 H

D